

Dectiver® -F

Endectocide and broad spectrum Fasciolicide (Injectable solution of Ivermectin and Clorsulon)

REG. SAGARPA Q-2083-009



FORMULA

Each mL contains:

Ivermectin 10 mg

Clorsulon 100 mg

Vehicle cbp 1 mL

PRESENTATIONS: Plastic bottles of 500 mL

DESCRIPTION

DECTIVER® -F is a sterile injectable solution of Ivermectin and Clorsulon ready to be used. This indicated in the treatment of internal parasites and including the fasciola in both its larval stage as an adult and as a coadjutant in the control of the external parasites of cattle and sheep so which helps improve your conversion efficiency food and weight gain contributing to have a greater weight to market. It is administered by subcutaneously at neck level or in front of the palette

INDICATIONS

DECTIVER® -F is effective in the treatment and control of adult and larval stages of internal parasites for up to 28 days depending on parasite including the fasciola as well as the external parasites in cattle.

DECTIVER® -F acts in the effective control of the following parasites of cattle and sheep:

Ostertagia ostertagi (includes hypobiotic stage)

Ostertagia spp

Bunostomun phlebotomun

Cooperia spp

Haemonchus spp

Oesophagostomum spp

Trichostrongylus spp

Nematodirus spp

Toxocara vitulorum

Mecistocirrus digitatus

Strongyloides papillosus

Ovine chabertia

Trichuris ovis

Dictyocaulus spp

Bovine parafilaria

Thelazia spp

Protostrongylus rufescens

Fasciola:

Hepatic Fasciola

Giant Fasciola

Control the following external parasites :

Fly larvae: Hipoderma bovis

H. lineatum

Dermatobia hominis

Oestrus ovis

Chrysomya bezziana

Mite producers of scabies:

Psoroptes ovis var bovis

P. communis var ovis

Choriotes bovis

Against lice:
Haematopinus eurysternus
Solenopotes capillatus

Assistant in the control of :
Damalinia bovis

ROUTE OF ADMINISTRATION

The only recommended route of administration of DECTIVER® -F is the subcutaneous route at the neck or in front of the paddle on the sheep in the axillary region. DECTIVER® -F can be administered with a syringe of automatic or conventional type.

DOSE

Cattle and sheep: 1 mL per 50 kg of weight alive. WARNINGS: It should not be applied intramuscularly or intravenous Keep out of reach of children. Stay in a cool place and do not expose yourself to Sun rays.

This product must not be used 28 days before the sacrifice of animals destined for consumption human. Do not use milk for human consumption of animals treated up to 28 days after the last application. Aquatic organisms are highly sensitive to the toxicity of Ivermectin, which is why empty containers and any residual content they must be disposed of safely Burying or incinerating them

VETERINARY USE.

CONSULT THE VETERINARY PHYSICIAN.

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION