

## **Dectiver® TF**

Oral Suspension of Ivermectin and Triclabendazole

REG. SAGARPA Q- 2083-170



### **FORMULA**

Each mL contains: Ivermectin 2 mg

Triclabendazole 120 mg

Vehicle cbp 1 mL

### **PRESENTATIONS**

1 liter plastic jars and 5 liter jugs.

### **INDICATIONS**

Desparasitante of wide spectrum, whose active ingredients triclabendazole and ivermectin are indicated for the control of Fasciola hepatica both in the juvenile and adult phases as well as in the control of parasite infections gastrointestinal and pulmonary diseases such as Ostertagia, Haemonchus, Trichostrongylus, Cooperia, Oesophagostomum, Trichuris, Strongyloides, Nematodirus, Chabertia, Dactylocaulus, Oestrus (larval stage), sucking lice

### **USE**

Cattle, sheep and goats.

### **DOSAGE AND ROUTE OF ADMINISTRATION**

Cattle: 5 mL per 50 kg of live weight.

Sheep and goats: 0.5 mL per 5 kg of live weight.

### **WARNINGS**

This product must not be used 28 days before the slaughter of animals destined for consumption human. Sacrifice the animals for human consumption after 21 days of the last treatment. When the milk or by-products of the milk are for human consumption, it should not be administered to cows 28 days before calving or during lactation. Aquatic organisms are highly sensitive to the toxicity of Ivermectin, which is why empty containers and any residual contents should be disposed of safely burying or incinerating them. Keep out of reach of children. Stay in a cool place and do not expose yourself to sunlight. Shake well before use.

CONSULT A VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION