

# Diramox® Soluble

Long-acting injectable amoxicillin

## REG. SAGARPA Q-2083-133

### FORMULA

Each 100 g contain:

Amoxicillin (as Amoxicillin trihydrate) 75 g Excipient, c.b. 100 g

### PRESENTATIONS

Jar with 900 g.

### INDICATIONS

Indicated for the treatment of bacterial infections susceptible to Amoxicillin in birds and pigs. As well as adjuvant in infections secondary to a viral disease.

### DOSE

Treated animals should not have access to another source of water during treatment. Administer 15 to 20 mg of active ingredient per kg of body weight.

Chickens: Dissolve 100 g in 450 l of fresh drinking water. This dosage will provide medication to 5,000 kg of weight, for a day. Administer medicated drinking water every third day (two days of medication). In severe cases, administer every third day for 5 days (3 days of medication).

Ducks: Administer 20 mg / kg of water in drinking water. The administration should be every third day, for three days (two days of medication). Because the water that is ingested varies depending on the temperature, light and diet, the exact dilution can not be provided. In any case, the quantity of Diramox® Soluble must be calculated taking into account the water ingested and the total weight of the ducks involved.

Pigs: Administer 20 mg / kg of water daily in drinking water. The dose should be divided and administered at approximate intervals of 12 hours for 5 days.

Turkeys: Administer in drinking water 15 to 20 mg Amoxicillin per kg of weight. The administration must be provided during 5 days in an alternate way (3 days of medication). The dosage should be calculated taking into account the normal amount of water to be ingested and the total weight of the turkeys involved.

### ROUTE OF ADMINISTRATION

Oral.

### WARNINGS

- Do not administer to animals with an allergic background to penicillin.
- During the treatment, the animals should not be destined for human consumption.

- Period of withdrawal:

Chickens: 24 hours.

Turkeys: 5 days.

Ducks: 7 days.

Pigs: 24 hours.

- Do not consume the eggs of treated birds until 24 hours after the last treatment.



CONSULT YOUR VETERINARY PHYSICIAN YOUR SALE REQUIRES MEDICAL RECIPE