

Dorso Pet®

permethrin

INSECTICIDE OF LIVESTOCK USE

POUR ON / OIL LIQUID

For the control of Fleas, Lice and Ticks.

RSCO-PEC-INAC-0156-395-378-007



PERCENTAGE COMPOSITION:

PERMETHRINE: 3-phenoxybenzyl (1RS, 3RS, 1RS, 3SR) -3- (2,2-dichlorovinyl) -2,2-dimethylcyclopropanecarboxylate (60% trans isomer and 40% cis isomer) 7.4% (Equivalent to 64.9 g from ia / L to 20°C)

INERT INGREDIENTS: Synergistic agent and diluent 92.60%

TOTAL 100%

Net Content:

1 package with 3 pipettes each, 0.5, 1, 2 and 3 mL

INDICATIONS

DORSO PET® is a product based on a pyrethroid and a synergist, with action on populations of fleas, lice and ticks.

DORSO PET® controls new infestations by fleas for up to 28 days after the application of 1 pipette.

For the treatment and control of allergic dermatitis by flea bite (DAP), monthly applications are recommended in allergic animals and other dogs that live with them.

DORSO PET® controls infestations caused by chewing lice of the genus *Trichodectes canis* in dogs and puppies.

DORSO PET® controls reinfestations caused by all stages of development of ticks of the genus *Rhipicephalus*, *Dermacentor* and *Ixodes*.

MECHANISM OF ACTION

Permethrin is a pyrethroid that has neurotoxic action on the nerve membranes of ectoparasites, causing excitation, coordination disorders, paralysis and death. The synergistic effect of piperonyl butoxide on the resistance enzymes prevents the degradation of permethrin by multiplying its action.

ADMINISTRATION AND MODE OF USE

DOSAGE: DORSO PET® are single-dose pipettes, each pipette contains the minimum recommended dose for treatment with Permethrin in dogs:

A 0.5 mL pipette for dogs weighing less than 5 kg

One 1 mL pipette for dogs weighing between 6 - 15 kg

A 2 mL pipette for dogs weighing between 16 - 30 kg

A pipette of 3 mL for dogs weighing between 31 - 45 kg

DORSO PET® can be applied in puppies from 8 weeks of age, as well as pregnant or lactating females.

Mode of use: Topical use only. Apply a pipette every 28 days for as long as the Veterinarian advises.

Avoid contact with the dog's skin after the application of DORSO PET® until the product is absorbed into the skin. Do not bathe the dog two days before or two days after treatment.

DORSO PET® is toxic to cats.