

Estreptopen®

Blend of penicillins and Streptomycins
REG. SAGARPA Q-2083-003

FORMULA

Each 1'000,000 UI contains:
Penicillin G Procain 375 000 IU
Penicillin G Sodium Penicillin G 125 000 IU
Dibencllic 500 000 IU
Streptomycin (as Streptomycin Sulfate) 375 mg
Dihydrostreptomycin (as dihydroestepotomycin sulfate 375 mg

PRESENTATIONS

Bottles with antibiotic powder of 2, 4, 12 and 50 million IU with its diluent respective.

INDICATIONS

Combination of:
Penicillins of short and long duration that act against gram-positive bacteria
Streptomycin
Dihydrostreptomycin that act against gram negative bacteria, which gives it a broad spectrum of action.

It is indicated in all species as initial therapy in various systemic infections. The combination penicillin-streptomycin is effective in various mixed infections such as middle ear infections and infections of the navel, as well as diseases of the respiratory system.

The combination penicillin-streptomycin is indicated as a complement to other therapies to combat secondary infections that commonly accompany fever and mastitis.

Due to its broad spectrum of action is useful in the treatment of infections in newborn animals.

SPECIES FOR WHICH THE PRODUCT IS RECOMMENDED

Bovines, swine, horses and small species.

DOSE, ROUTE OF ADMINISTRATION AND INSTANCES OF USE

Apply from 11,000 to 22,000 IU per kilogram of live weight.

Mix perfectly with the accompanying diluent.

Administer by deep intramuscular injection only. For best results:

- Trim the hair at the application site, wash with soap and water disinfect with iodine stain, alcohol or some another local germicide.
- Use sterile syringes and needles. Use 16 or 18 gauge for large animals and 19 or 20 for animals little ones.
- Mix the powder perfectly with the accompanying diluent.
- Fill the syringe with the amount of product needed.
- Insert the needle into the injection site with negative pressure to make sure that the needle is not in a blood vessel and then inject the product into the muscle.



WARNINGS

- There may be cases of hypersensitivity in individuals allergic to penicillin. In this case, the treatment should be discontinued and the subject should be treated with the agents usually used (amines vasopressors, corticosteroids, and antihistamines).
- Do not administer to animals with a history of allergic reactions or hypersensitivity to penicillins or cephalosporins or their analogues.
- To avoid discomfort after the application, it is recommended not to administer more than 10 mL in the same injection site. Store in a cool place, protect yourself from sunlight.
- Keep out of the reach of children.

RETIREMENT PERIOD

This product should not be used 30 days before the slaughter of animals destined for consumption human.

Milk from treated cows should not be used until 72 hours after treatment.

CONSULT THE VETERINARY PHYSICIAN.

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION.