

# Lapimicina® LA

Long-acting oxytetracycline  
REG.SAGARPAQ-2083-051



## FORMULA

Each mL contains: Oxytetracycline base 200 mg (as Oxytetracycline hydrochloride)  
Vehicle cbp 1 mL

## PRESENTATION OF THE PRODUCT

Bottle with 100, 250 and 500 mL.

## INDICATIONS

Oxytetracycline with long action, indicated in the treatment of pneumonia, diarrhea, pododermatitis, keratoconjunctivitis, actinobacillosis, leptospirosis, anaplasmosis, mastitis, metritis and agalactea as well as infectious septicemic processes.

## SUSCEPTIBLE ORGANISMS

Pasteurella spp., Haemophilus spp., Moraxella bovis, Fusobacterium necrophorum, Escherichia coli, Actinobacillus lignieresii, Leptospira spp., Staphylococcus spp., Streptococcus spp., Actinobacillus pleuropneumoniae

## USE

Bovines, pigs, sheep and goats.

## DOSAGE, ADMINISTRATION AND INSTRUCTIONS

Administer by deep intramuscular injection at a rate of 20 mg of Oxytetracycline per kg of weight. Equivalent to apply 1 mL of product for every 10 kg of weight. Do not apply more than 20 mL per injection site. In animals of less than 10 kg in weight, recommends the subcutaneous application of the product.

## INTERACTIONS AND INCOMPATIBILITY WITH OTHER PRODUCTS

Bacteriostatic drugs can interfere with the bactericidal action of penicillin so that Advise to avoid giving treatment with Lapimicina® LA in conjunction with penicillins.

## PRECAUTIONS

Overdose may cause the residues of the antibiotic to remain beyond the period of suspension. There have been allergic or anaphylactic reactions, sometimes fatal, in hypersensitive animals after an oxytetracycline injection. If this occurs, discontinue the treatment and consider the administration of antihistamines. The use of the product may cause irritation local transient manifested by inflammation and discoloration at the injection site. The pain produced by the injection can be minimized by heating the solution at body temperature before the administration and avoiding a bad injection technique. After treatment, animals can present transient hemoglobinuria. Like all antibiotic preparations, the use of this product They can result in an overgrowth of a resistant organism, including fungi. The absence of a favorable response after treatment, or the development of new signs or symptoms, may suggest the overgrowth of non-susceptible organisms. If superinfection occurs, the use of this product should be discontinued and the treatment. Do not use for human consumption, the milk of treated animals until 60 hours after the last application. Store in a cool, dry place protected from sunlight. Keep out of reach of children.

## RETIREMENT PERIOD

This product should not be used 30 days before the slaughter of animals destined for human consumption.

## CONSULT A VETERINARY

**YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION**