

"AGRICULTURAL USE"



# **MANAGER<sup>®</sup>** 350sc **IMIDACLOPRID**

Imidacloprid Insecticide,  
Concentrated Suspension

**PERCENTAGE COMPOSITION**

**% in weigh**

**Active ingredient:**

**32.65**

**Imidacloprid:**

(E) -1- (6-Chloro-3-pyridylmethyl) -N-nitroimidazolidin-2-ylideneamine (Equivalent to 350 g i.a./L at 22 ° C)

**Inert ingredients:**

**67.35**

Emulsifier, solvent and vehicle

**Total**

**100.00**

**Registry: RSCO-INAC-0199-339-064-033**



"Harmful if swallowed"  
"Harmful if inhaled"

Holder of the registration, Formulated and  
Distributed by: LAPISA, S.A. DE C.V.  
Carr. La Piedad-Guadalajara Km. 5.5  
Col. Camelinas, C.P. 59375,  
La Piedad, Michoacán, Mexico.  
Tel. : 01 (352) 526 13 00.  
[www.lapisa.com](http://www.lapisa.com)  
MADE IN MEXICO

"THIS MATERIAL IS NOT A LABEL, THE FUNCTION OF THIS DOCUMENT IS ILLUSTRATIVE."

# CAUTION

## "STOP: READ THE LABEL BEFORE USING THE PRODUCT"

### "PRECAUTIONS AND WARNINGS FOR USE"

Use the appropriate and complete protective equipment necessary to safely perform the handling operations for the preparation, the loading of the application equipment and its application: cotton cap and overalls, face mask against liquids, goggles or chemical-type lenses, gloves and Neoprene boots.

#### "PRECAUTIONS DURING PRODUCT HANDLING":

"DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT".

"DO NOT TRANSPORT OR STORE ALONG WITH FOOD PRODUCTS, CLOTHES OR FORAGE."

"KEEP OUT OF REACH OF CHILDREN, PREGNANT WOMEN, BREASTFEEDING AND DOMESTIC ANIMALS".

"WASH YOUR HANDS AFTER USING THE PRODUCT AND BEFORE CONSUMING FOOD".

"DO NOT STORE IN HOUSES ROOM".

"DO NOT USE THE PRODUCT WITHOUT THE PROPER PROTECTION EQUIPMENT".

"PERSONS UNDER 18 YEARS OLD SHOULD NOT HANDLE THIS PRODUCT".

"AT THE END OF THE WORK DAY BATH AND PUT CLEAN CLOTHES".

"DO NOT INGIRE, INHALE AND AVOID CONTACT WITH EYES AND SKIN".

"DO NOT MIX OR PREPARE THE PRODUCT WITH YOUR HANDS, USE GLOVES, USE A PIECE OF WOOD OR OTHER APPROPRIATE MATERIAL".

"DO NOT UNLOCK THE NOZZLE WITH THE MOUTH".

### "IN CASE OF INTOXICATION, TAKE THE PATIENT WITH THE DOCTOR AND SHOW THIS LABEL"

#### "FIRST AID"

In case of intoxication get medical attention immediately or call the Poison Attention Center (ATOX) for medical assistance 24 hours a day at 01-800-000-2869, meanwhile, it is recommended to apply the following first aid measures:

**Inhalation:** Move contaminated person to fresh air. If you do not breathe, give artificial respiration, preferably mouth to mouth.

**Ingestion:** Immediately give one or two glasses of water and induce vomiting by gently touching the back of the throat with a finger, or dull object. Do not induce vomiting or administer anything by mouth to an unconscious person.

**Skin:** Wash with plenty of soap and water.

**Eyes:** Rinse with plenty of water for at least 15 minutes.

### "RECOMMENDATIONS TO THE DOCTOR"

**Chemical Group:** Cloronicotiniles.

**Antidote:** No specific antidote is known. Use symptomatic treatment.

**Treatment:** Therapy in children should be supervised by a doctor. Check blood pressure and pulse rate, give artificial respiration if signs of paralysis appear.

Apply therapeutic measures for elimination of the substance of the body (gastric lavage, saline laxatives).

Signs and Symptoms of Poisoning:

**Oral:** diarrhea and salivation. **Dermal:** tingling that disappears after a short time, shuddering and hypersensitivity. **Inhalation:** apathy or impassivity, muscular depression, respiratory disturbances, tremor and in severe cases muscle cramps.

### ECOTOXICOLOGICAL CATEGORY:

THIS PRODUCT IS HIGHLY TOXIC FOR LIVER BLOODY ANIMALS (BIRDS).

THIS PRODUCT IS EXTREMELY TOXIC FOR ANIMALS (AQUATIC INVERTEBRATES).

THIS PRODUCT IS HIGHLY TOXIC FOR BEES. DO NOT APPLY WHEN THE CULTIVATION OR WEEDS ARE IN FLOWER AND WHEN THE BEES ARE RELEASED.



**“ÚSESE EXCLUSIVAMENTE EN LOS CULTIVOS Y PLAGAS AQUÍ RECOMENDADOS”**  
**“INSTRUCCIONES DE USO”**  
**“SIEMPRE CALIBRE SU EQUIPO DE APLICACIÓN”**

CROP	PLAGUE	DOSE
Pumpkin, Zucchini, Melon, Cucumber, Watermelon (21)	White fly ( <i>Bemisia tabaci</i> )	0.75 – 1.0 L/Ha
Avocado (14)	White fly ( <i>Paraleyrodes perseae</i> ) Trips ( <i>Scirtothrips perseae</i> )	300 – 600 mL/Ha 450-600 mL/Ha
Tomato, Chile, Eggplant (21)	White fly ( <i>Bemisia tabaci</i> ) Aphids ( <i>Myzus persicae</i> )	0.75 – 1.0 L/Ha
Lettuce (21)	Aphids ( <i>Myzus persicae</i> ) Trips ( <i>Frankliniella occidentalis</i> )	160-200 mL/Ha
Broccoli, Cabbage, Cauliflower, Brussels sprouts (21)	Ashen aphid ( <i>Brevicoryne brassicae</i> )	160-200 mL/Ha
Vine(21)	Floury louse ( <i>Planococcus ficus</i> )	1.0-1.5 L/Ha

( ) Security interval in days between the last application and harvest.

**TIME OF REENTRANCE TO THE TREATED ZONES: DO NOT ENTER THE TREATED AREA BEFORE 24 HRS.**

**METHOD FOR PREPARING THE PRODUCT**

Calibrate the application equipment, in a separate container place a small amount of water, add the product and shake, then empty this premix to the equipment tank and add all the water calculated in the calibration.

**Avocado:** Perform two applications to the foliage at 14-day intervals. Time of application in flowering and fruiting, with a volume of application: 1000 L of water / Ha.

**Tomato, chili and eggplant:** Make a drench application to the neck of the plant, 15 days after the transplant, with an application volume of 12.5 mL of the mixture per plant.

**Lettuce:** Make an application to the foliage, with a water application volume of 362 to 462 L / Ha.

**Broccoli, Cabbage, Cauliflower, Brussels Cabbage:** Make an application to the foliage, with a volume of water application of 369 to 469 L / Ha.

**Vine:** Perform a drench application to the base of the plant, with an application volume of 250 mL of the mixture per plant.

**CONTRAINDICATIONS:** Do not apply with strong winds or temperatures above 35 ° C.

**PHYTO-TOXICITY:** This product is not phytotoxic in the crops indicated here if it is applied at the doses recommended in this label.

**INCOMPATIBILITY:** Do not mix Imidacloprid with strong alkaline reaction products.

**RESISTANCE MANAGEMENT**

"TO PREVENT THE DEVELOPMENT OF RESISTANT POPULATIONS, ALWAYS RESPECT THE DOSE AND APPLICATION FREQUENCIES; AVOID THE REPEATED USE OF THIS PRODUCT, ALTERING IT WITH OTHER CHEMICAL GROUPS OF DIFFERENT MODES OF ACTION AND DIFFERENT MECHANISMS OF DETOXIFICATION AND THROUGH THE SUPPORT OF OTHER CONTROL METHODS ".

**WARRANTY:** LAPISA S.A. DE C.V., guarantees the composition of the product in accordance with what is specified in the present label. However, for no reason assumes responsibility for the misuse of the product or for any damage that could be caused.

