

BOVIMUNE MASTER L5®

REG. SAGARPA B-2083-022



FORMULA

Each dose of 3 mL contains:

Freeze-dried fraction: Live modified parainfluenza virus (PI3) and Bovine Syncytial Respiratory Virus (BRSV); Excipient c.b.p. 1 dose
Liquid fraction: Inactivated Bovine Infectious Rhinotracheitis (IBR) Virus, Bovine Viral Diarrhea (DVB) types I and II; and inactivated bacteria of *Leptospira canicola*, *L. grippotyphosa*, *L. hardjo*, *L. icterohaemorrhagiae*, and *L. pomona*.
Excipient c.b.p. 3 mL

PRESENTATIONS

Bottle with lyophilized:
10 and 25 Dosage.
Sterile liquid fraction:
30 and 75 mL.

INDICATIONS

It is indicated for the vaccination of dairy cattle or healthy fattening, including pregnant cows; for the prevention of infections caused by Infectious Bovine Rhinotracheitis, Bovine Viral Diarrhea types I and II, Parainfluenza3 and Respiratory Syncytial Bovine, as well as infections caused by the mentioned serotypes of *Leptospira*.

INDICATED SPECIES

Bovine.

DOSE

Reconstitute the lyophilized fraction with the liquid fraction and shake well.
Apply aseptic technique 3 mL in healthy animals.
You can repeat the dose 14 to 28 days after the first application.
Annual vaccination is recommended.

ADMINISTRATION MODE

Intramuscular or subcutaneous.

WARNINGS

Once the container is opened, use the entire contents.
Keep refrigerated at a temperature between 2° and 7° C, do not freeze.

Any biologic can cause anaphylactic reactions. In such case, the use of epinephrine followed by supportive therapy is recommended.

Do not vaccinate 21 days before the slaughter of animals intended for human consumption. Dispose of the container securely, lock or incinerate.

It contains Gentamicin as a preservative.

CONSULT THE VETERINARY PHYSICIAN

Produced by:
Diamond Animal Health, Inc.
Des Moines, Iowa 50327 U.S.A.
Veterinary License
in the USA No. 213.