

Lapisa **Minoxel[®] L.A.** Cerdos

SUSPENSION OLEOSA INJECTABLE

REG SAGARPA Q-2083-199 VETERINARY USE

FORMULA

Each mL contains: Ceftiofur Crystalline Free Acid equivalent to Ceftiofur 100 mg
Vehicle cbp1 mL



INDICATIONS

Ceftiofur Crystalline Free Acid (Lapisa Minoxel[®] LA Pigs), is indicated for the treatment of Porcine Respiratory Disease associated with Actinobacillus pleuropneumoniae, Pasteurella multocida, Haemophilus parasuis and Streptococcus suis .

DESCRIPTION OF THE DISEASE

Respiratory-type disease caused by Actinobacillus pleuropneumoniae, Pasteurella multocida, Haemophilus parasuis and Streptococcus suis that affects the respiratory tract of the pig. The transmission is from pig to pig, the high densities and the practices of inadequate management increase the predisposition to the disease, the signs can vary from a severe respiratory disease of the type chronic until sudden death. The most critical moment is in growing animals, because the pigs are regrouped and mixed, suffer from high levels of stress, lose passive immunity and result in cases of pneumonia two or three weeks after weaning.

DOSE

A single application of 5 mg of Ceftiofur per kg of body weight (1 mL per 20 kg of weight). If necessary, you can repeat the dose at 7 days after the first application.

ROUTE OF ADMINISTRATION

Intramuscular on the neck table..

WARNINGS

Keep out of reach of children and pets. Avoid contact with the skin of people allergic to β -lactams. Can thrombocytopenia and reversible anemia occur in animals given three to five times the recommended dose. Pigs weighing more than 40 kg will require more than one injection as they can not exceed 2 mL per application site. This product should not be used 14 days before the slaughter of animals destined for human consumption.

CONTRAINDICATIONS

Do not administer to animals that have previously shown hypersensitivity to Ceftiofur or other β -lactam antibiotics. People who handle this medication should be careful to avoid repeated or prolonged exposure to prevent sensitization to antimicrobials that may occur in susceptible individuals. Avoid direct contact with eyes, mouth or skin. Those with known hypersensitivity should avoid exposure to this medication.

PRESENTATION

100 mL

STORAGE

Store at controlled room temperature between 20 and 25 °C.

CONSULT A VETERINARY

YOUR SALE REQUIRES A MEDICAL RECIPE