

OVOSURE®

Reg. SAGARPA Q-2083-163

FORMULA

Each ml contains: Gonadorelin acetate 100 mcg.
Vehicle, cbp 1 mL

PRESENTATIONS

Flasks with 50 mL.



INDICATIONS

Ovasure ® is indicated in the treatment of ovarian cysts, prevention of faults in the ovulation, induces ovulation improving postpartum fertility

MECHANISM OF ACTION

It is a solution based on gonadorelin acetate, a synthetic analogue of the hormone GnRH gonadotropins; Its function is the production and secretion of luteinizing hormone (LH) and follicle stimulating hormone (FSH). Both LH and FSH act in the ovary, FSH stimulates the growth, development and function of the follicle; LH induces ovulation, in addition to the development and function of Corpus luteum. Gonadorelin is rapidly absorbed once administered intramuscularly, after administration of absorption, has a rapid distribution reaching therapeutic levels at the site of action. The plasma half-life of gonadorelin is 20 minutes, it is metabolized by enzymes peptidases, the main route of elimination of metabolites is urine

USE

Bovine

DOSAGE AND ROUTE OF ADMINISTRATION

Cattle: Treatment of ovarian cysts: 5.0 mL.

Ovulation delay: 2.5 mL.

Improvement of postpartum fertility: 2.5 mL.

Stroke synchronization: 1.0 mL.

WARNINGS

Withdrawal time:

Meat: 0 days.

Milk: 0 days.

Store at temperatures below 25 ° C, in a dry place protected from light.

Keep out of reach of children.

CONSULT THE VETERINARY

REQUIRE MEDICAL RECEIPT

EXCLUSIVE USE OF THE VETERINARY