BOVIMUNE PROTECTOR L5®

REG. SAGARPA B-2083-023

FORMULA
Each dose of 2 mL contains:
Lyophilized fraction: Live modified viruses of Bovine Infectious Rhinotracheitis (IBR), Bovine Viral Diarrhea (DVB) type I and type II, Bovine Syncytial Respiratory Virus (BRSV) and Parainfluenza3 (PI3).
Diluente: Inactivated bacteria of Leptospira canicola, L.grippotyphosa, L.hardjo, L. icterohaemorrhagiae, and L. pomona.
Excipient c.b.p. 2 mL.

PRESENTATIONS
Bottle with lyophilized:
10 Dosage.
Bottle with sterile diluent:
20 mL.

INDICATIONS
Live modified vaccine for the prevention of infections caused by bovine infectious rhinotracheitis viruses, Bovine Viral Diarrhea types I and II, Bovine Syncytial Respiratory Virus and Parainfluenza3 in healthy cattle, as well as infections caused by the mentioned serotypes of Leptospira.

INDICATED SPECIES
Bovine.

DOSE
Transfer aseptically the entire contents of the diluent bottle to the bottle containing the lyophilized fraction and shake well. Apply using aseptic technique 2 mL in healthy animals. Calves vaccinated before 6 months should be re-vaccinated at six months of age or at the time of weaning.
Annual vaccination is recommended.

ADMINISTRATION MODE
Intramuscular or subcutaneous.

WARNINGS
Do not use in pregnant females since it can cause abortion, nor apply in calves that breastfeed.
When reconstituting the vaccine do not use syringes or needles disinfected with chemical agents.

Once the container is opened, use the entire contents.
Keep refrigerated at a temperature between 2° and 7° C, do not freeze.
Do not expose the product to sunlight.
Any biologic can cause anaphylactic reactions. In this case, the use of epinephrine followed by appropriate support therapy is recommended.
Do not vaccinate 21 days before the slaughter of animals intended for human consumption. Dispose of the container securely, lock or incinerate.

CONSULT THE VETERINARY PHYSICIAN

Produced by:
Diamond Animal Health, Inc.
Des Moines, Iowa 50327 U.S.A.
Veterinary License
in the USA No. 213.