

# TICOFF®

cypermethrin

GARRAPATICIDA AND MOSQUICIDA FOR LIVESTOCK, EMULSIONABLE CONCENTRATE

REG. RACO INAC 111-12-2083-001

## FORMULA

### Percentage composition

Cypermethrin: 2 cyano-3 Phenoxybenzyl, Trans 3- (2,2-dichlorovinyl -3,2)

Propane carboxylate cycle.

(Equivalent to 200 g of i.a./L at 20 ° C)

Emulsifier, solvent

Total

### By weight

20.00%

80%

100%



## PRESENTATION:

30 mL.

## PHARMACOLOGICAL PROPERTIES

Cypermethrin is a synthetic pyrethroid used as an insecticide for the control of infinity of pests, directly on animals or crops, including cotton moths, fruits, and vegetable harvest. It is also used in cracks and crevices for the control of insects in stores, warehouses, industries, houses, apartments, nurseries, laboratories, ships, trains, buses, trucks, and airplanes. It can also be used in schools (except food areas) hospitals, hotels, food processing plants and as a repellent barrier treatment in horses.

When applied to animals, it is absorbed by skin in a low proportion (1% in 24 hours in rats), although its detoxification is very rapid and does not accumulate in animal tissues or in the environment, unlike pesticides. It is rapidly metabolized in the liver by hydrolysis and oxidation reactions

## USE RECOMMENDED BY THE MANUFACTURER

Ticoff® acts on all stages of the ticks of one or several hosts with an excellent efficacy and prolonged residual effect (10-14 days). Ticoff® also controls other external parasites that affect cattle causing irritation, pain, and discomfort, such as: stable fly, paddle fly, face fly, as well as horseflies, lice and mites.

## SPECIES FOR WHICH THE PRODUCT IS RECOMMENDED

Cattle.

## DOSAGE AND ADMINISTRATION MODE

Immersion bath:

Load:

Cubicize the volume of the bath, use 1 liter of Ticoff ®.

Place the volume of the bath, use 1 liter of Ticoff ®.

Place the volume of the bath, use 1 liter of Ticoff ® per each for every 1000 liters of water. Pre-mix the required amount of Ticoff® in an equal amount of water and add this mixture to the water of the immersion bath.

Recharge:

Add 1.5 liters per 1000 liters of water.

Spray bath:

Prepare a mixture with 15 ml of Ticoff ®

Prepare a mixture with 15 ml of Ticoff ®

Prepare a mixture with 15 ml of Ticoff ® for every 10 liters of water per 10 liters per 10 liters of water and sprinkle with fine drop to the whole animal.

Facilities:

Apply 3 mL of Ticoff ® per liter of water.

Apply 3 mL of Ticoff ® per liter of water

Apply 3 mL of Ticoff ® per liter of water.

## RETIREMENT PERIOD

Immersion baths: Animals of

Meat: 4 days Milk: 1 day

## •PRECAUTIONS DURING PRODUCT HANDLING

- DO NOT EAT, DRINK OR SMOKE WHEN YOU ARE USING THIS PRODUCT.
- NO SE TRANSPORTE NI ALMACENE JUNTO A PRODUCTOS ALIMENTICIOS, ROPA O FORRAJES.
- MANTENGA LA FUERA DEL ALCANCE DE LOS NIÑOS, MUJERES EMBARAZADAS, EN LACTANCIA Y ANIMALES DOMÉSTICOS.
- LÁVESE LAS MANOS DESPUÉS DE UTILIZAR EL PRODUCTO Y ANTES DE CONSUMIR ALIMENTOS.
- SIN USO EL PRODUCTO SIN EL EQUIPO DE PROTECCIÓN ADECUADO.
- NO SE INGIERA, INHALA Y EVITE SU CONTACTO CON LOS OJOS Y PIEL.
- NO MEZCLE O PREPARE EL PRODUCTO CON LAS MANOS, USE GUANTES, USE UN TROZO DE MADERA U OTRO MATERIAL APROPIADO.
- VETERINARY USE
- CONSULT YOUR PHYSICIAN BEFORE USING THE VETERINARY THIS PRODUCT