

Antilmin 12%

Injectable anthelmintic

REG. SAGARPA Q-2083-034

PRESENTATIONS

500 mL.

FORMULA

Each 100 mL contains:

Levamisole Hydrochloride 12.0 g

(equivalent to levamisole base 10.2 g) Vehicle

c.b.p. 100 mL

INDICATIONS

Broad spectrum anthelmintic for the treatment of pulmonary and gastrointestinal parasitosis, both in mature and immature forms in cattle, sheep, goats and pigs.

INDICATED SPECIES

Bovine, ovine, porcine and goat.

DOSE

Gastrointestinal parasites 1 ml per 20 kg of live weight.

Pulmonary parasites 1 ml per 15 kg of live weight. Maximum dose 15 ml per application site.

ROUTE OF ADMINISTRATION:

Deep intramuscular.

WARNINGS

Consult a veterinarian before using the product on highly weakened animals. Dosage should be monitored by an accurate estimate of the animal's weight

Do not use this product 3 days before the slaughter of animals destined for human consumption.

Do not use the milk of cows treated with this product until 72 hours after the last application.

In some animals excessive salivation or the formation of "foam" in the snout can be observed in a transitory form which will disappear by itself in a few hours.

Store in a cool, dry place protected from sunlight.

Keep out of the reach of children.

CONSULT THE VETERINARY PHYSICIAN.

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION

