

Bromucol®

Mucolytic and expectorant.

REG. SAGARPA Q-2083-154

FORMULA

Each mL contains: Bromhexine Hydrochloride 3 mg
Vehicle, c.b.p. 1 mL



INDICATIONS

Mucolytic and adjuvant expectorant for the treatment of respiratory tract diseases that occur with mucus, increased viscosity and cases of retention of secretions, such as chronic pneumonias, acute and chronic bronchopneumonia, acute and chronic bronchitis, pharyngitis, laryngitis, tracheitis and sinusitis. Also in the presence of simple processes such as cough and catarrh.

DOSE

0.5 mg of Bromhexine Hydrochloride per kg of body weight, equivalent to 1 mL per 6 kg of body weight.

In general, it is recommended to apply the following doses:

Bovine and equine adults: 25 to 30 mL.

Calves and calves: 8 to 20 mL.

Pigs: 5 to 15 mL.

Piglets: 2 to 3 mL.

Sheep: 5 to 10 mL.

Dogs: 2 to 5 mL.

Cats: 1 to 2 mL.

TREATMENT DURATION

5 days.

ROUTE OF ADMINISTRATION

Intramuscular and slow intravenous.

MANUFACTURER'S RECOMMENDATIONS:

To solve respiratory problems, it is recommended to make a mixture of Bromucol® with Minoxel g in the following way:

100 mL of Bromucol® a bottle of Minoxel 4 g + 100 mL of sterile water and shake, for a total volume of 200 mL.

Apply 1 mL per 10 kg in pigs for the treatment and control of porcine bacterial respiratory disease (porcine bacterial pneumonia) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Salmonella choleraesuis* and *Streptococcus suis*.

Once reconstituted, store in refrigeration for no more than 7 days.

WARNINGS

The meat and milk of treated bovines are not used for human consumption until 3 days after the last treatment.

The meat of treated pigs is not used for human consumption until 24 hours after the last treatment.

Keep out of reach of children. Store in a cool, dry place protected from light.

VETERINARY USE

CONSULT THE VETERINARY PHYSICIAN