

Coxizuril 5%

Prevention and control of Oral coccidiosis

Reg. SAGARPA Q-2083-155

FORMULA

Each mL contains:

Toltrazuril 50 mg

Excipient, c.b.p. 1 mL

PRESENTATIONS

Bottles with 250 mL.



INDICATIONS

It is indicated for the prevention and control of coccidiosis (*Isospora suis*) in piglets. In the piglets treated with COXIZURIL 5% there is a significant reduction of diarrheal cases in the second and third week of age, a better productive behavior is obtained and the use of antibiotics to control secondary infections decreases due to the notable reduction of cases of diarrhea.

Toltrazuril induces changes during the intracellular development stage of coccidia, mainly due to vacuolization of the endoplasmic reticulum, decreasing the nuclear division. COXIZURIL 5% acts against all stages of the development of coccidia, interferes with the division of the protozoan nucleus, affects the activity of the mitochondria and damages the corpuscles that form the wall in the microgametes. Therefore COXIZURIL 5% is an effective coccidicide since it destroys the parasites in the different stages of its development.

USE

Pigs.

DOSAGE AND ROUTE OF ADMINISTRATION

The recommended dose of toltrazuril in piglets is 20 mg / kg of live weight, which corresponds to 1 mL of COXIZURIL 5% per 2.5 kg of live weight. Apply a simple dose with the COXIZURIL 5% doser orally in piglets 3 to 5 days old.

WARNINGS

This product should not be used 70 days before slaughter of animals intended for human consumption.

Shake well before use.

Keep out of reach of children.

Store in a cool, dry place, protected from light.

CONSULT THE VETERINARY PHYSICIAN

YOUR SALE REQUIRES MEDICAL RECIPE VETERINARY USE