

# Diramox<sup>®</sup> Premix

Antibiotic Premix

**REG. SAGARPA Q-2083-081**

## FORMULA

Each kg. contains: Amoxicillin trihydrate 200 g  
Excipient, cbp 1000 g

## PRESENTATIONS

Bag of 20 kg



## INDICATIONS

Premix Antibiotic indicated for prevention from infections bacterial susceptible to amoxicillin in lactating birds, pigs and calves. As well as in infections secondary to a viral disease

## DOSE

Piglets: 1 - 1.5 kg. per ton of food for 5 - 7 days.

Sows: 1.5 - 2 kg. per ton of food for 5 - 7 days.

Birds: 1-2 kg. per ton of food for 5 - 7 days.

Breeding calves: 1.5 - 2.5 kg. per ton of food for 10 -20 days.

Weaning calves: 1,250 - 1,800 kg. per ton of food for 8-15 days.

Fattening: 4 - 6 kg. per ton of food during the first 15 - 21 days of the beginning of the cycle intensive fattening

## ROUTE OF ADMINISTRATION

Oral.

Mixed in the food.

## WARNINGS

Do not administer to animals with an allergic history to penicillin.

Do not use for human consumption the meat of treated animals until 48 hours after the treatment.

## CONSULT A VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION