

# Diramox® Premix 50%

Premezcla Antibiótica

REG. SAGARPA Q-2083-172



## FORMULA

Each kg contains: Amoxicillin trihydrate 500 g  
Excipient c.b.p. 1 kg

## PRESENTATIONS

Sacks of 10 kg.

## INDICATIONS

Antibiotic premix indicated for the prevention and treatment of bacterial diseases in birds, pigs and bovines; caused by Salmonella spp, Escherichia coli, Pasteurella spp, Mannheimia haemolytica, Histophilus spp, Haemophilus parasuis, Streptococcus spp, Actinobacillus spp and Clostridium spp.

## USE IN

Bovines, pigs and birds.

## DOSE

Species	Therapeutic Treatment
Poultry	400 to 800 g per ton of feed for 5 - 7 days.
Cattle	Breeding calves: 600 to 1000 g per ton of feed for 10 -20 days. Weaning calves: 500 to 720 g per ton of feed for 8 to 15 days. Calves fat: 1.6 to 2.4 kg per ton of feed during the first 15 - 21 days of the start of the intensive fattening cycle.
Swine	Piglets: 400 to 600 g per ton of feed for 5 - 7 days. Adults: 600 to 800 g per ton of food for 5 - 7 days.

## ROUTE OF ADMINISTRATION

Oral. Mixed with the food.

## WARNINGS

Withdrawal time:

Meat: calves and swine: 20 days.

Fattening birds: 7 days

Do not administer to animals with an allergic history to penicillin.

Do not administer this product to bovines in milk production that is intended for human consumption.

Do not administer this product to birds in egg production that is intended to human consumption.

Keep in a cool and dry place. Keep out of reach of children

CONSULT THE VETERINARY PHYSICIAN

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION