Electrolife

Soluble powder

Each 100 g contain:
Vitamin "C"
Vitamin "A"
Calcium chloride
Chloride potassium
Magnesium sulfate
Zinc sulfate
Manganese sulfate
Sodium Chloride
Excipient c.b.p.

0.100 g 50 000 U.I 2,000 g 3,400 g 0.700 g 0.380 g 0.430 g 57,200 g 100,000 g



PRESENTATIONS

Bag of 10 kg.

INDICATIONS

Coadjuvant in the treatment of dehydration, diarrhea, convalescence, stress or in all cases that rehydration is required.

(Transport, reception of cattle high temperatures, etc).

SPECIES FOR WHICH THE PRODUCT IS RECOMMENDED

Bovine, ovine, porcine and bird.

CONTRAINDICATIONS

Do not use this product in animals that are sensitive to the formula.

DOSAGE AND ADMINISTRATION MODE

Dissolve 250 grams of Electrolife in 200 L of drinking water, administering it for a period of 5 to 10 days.

ROUTE OF ADMINISTRATION

Oral dissolved in drinking water.

RETIREMENT PERIOD

0 days

SPECIAL STORAGE PRECAUTIONS

Store in a cool, dry place, protected from sunlight. Keep out of the reach of children.

CONSULT THE VETERINARY PHYSICIAN.