

Flunidin®

Analgesic, antipyretic and non-steroidal anti-inflammatory (Flunixin meglumine)
REG. SAGARPA Q-2083-074



FORMULA

Each mL contains:

Flunixin base 55 mg

(as Flunixin Meglumine USP) Vehicle, c.b.p.1 mL.

INDICATIONS

Cattle: In cattle FLUNIDIN® is indicated for the control of acute inflammatory processes associated with respiratory diseases such as the bovine respiratory complex (shipping fever).

Due to its antiendotoxic and antipyretic action, it is also indicated in septicemia, acute mastitis due to coliforms, as well as in the syndrome of the cow, and other inflammatory processes.

In any case, the primary cause of the inflammation must be determined and treated concomitantly (for example, by antibiotic therapy).

Swine: In pigs, FLUNIDIN® is indicated as adjuvant therapy for the Mastitis, Metritis, Agalactia (MMA) syndrome, and in the lactational failure of females caused by various inflammatory processes. Likewise as antipyretic in general.

Horses: In equines FLUNIDIN® is indicated for the relief of inflammation and pain associated with musculoskeletal disorders caused by unusual exercise (equine paralytic myoglobinuria, azothuria, Monday disease, stiff horses). It is also indicated for the relief of abdominal pain (colic). The use of FLUNIDIN® should be reserved for cases of colic that have been definitively diagnosed, otherwise the powerful analgesic effect may mask the deterioration of the general conditions of the animal, which could lead the clinician to a false sense of security.

USO

Bovinos, porcinos, equinos.

DOSAGE AND ROUTE OF ADMINISTRATION

Bovines, Swine, and Horses: Administer 2.0 mL / 50 kg of body weight.

Bovine and equine: Administer intravenously or intramuscularly. It is recommended to repeat the dose at a 12-hour interval.

Swine: Administer intramuscularly. It is recommended to repeat the dose with an interval of 12 hours.

WARNINGS

Animals should not be slaughtered for human consumption during treatment.

Do not use this product 24 hours before slaughter of animals intended for human consumption.

Do not use milk from animals treated with this product for human consumption, until 12 hours after the last application.

Store between 2 and 30°C (36 and 86°F).

Keep this and all medicines out of the reach of children.

It can be used with a wide margin of safety in cattle, swine and horses of any age and physiological condition since its toxic capacity is minimal. It can be used in conjunction with antimicrobials, however it is recommended not to mix Flunidin® with other medications in the same syringe.

PRESENTATIONS

Amber glass bottles with 100 and 250 mL.

CONSULT THE VETERINARY PHYSICIAN
YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION
VETERINARY USE