

Flupen

Association of antimicrobials with flumethasone
REG. SAGARPA Q-0402-101



FORMULA

Each million contains:

Penicillin G Procaine 750, 000 IU
Penicillin G Sodium 250, 000 IU
Dihydrostreptomycin Sulfate 0.500 g
Streptomycin Sulfate 0.500 g
Flumetasone 0. 312 mg

PRESENTATION

Ampla bottle with 2, 4 and 6 Millions.

INDICATIONS

Antimicrobial association with flumetasone, to fight infections that occur with inflammation that run with inflammation caused by Gram-positive and Gram-negative germs.

DOSE

Cattle and horses: 2 to 2.5 mL per 50 kg of weight.
Sheep and goats: 1 to 1.5 mL per 25 kg of weight.
Swine: 1 to 1.5 mL per 15 kg of weight. Canines and Felines: 0.5 to 2.0 mL according to weight.

ROUTE OF ADMINISTRATION

Deep intramuscular.

WARNINGS

Do not use this product 30 days before the slaughter of animals intended for human consumption.
Do not use milk from animals treated with this for human consumption product, up to 3 days after the last application
Store in a cool, dry place, protected from light.
Keep out of the reach of children

CONSULT THE VETERINARY DOCTOR
YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION