

Fluvin®

Anti-inflammatory glucocorticoid

REG. SAGARPA Q-2083-105

FÓRMULA

Each mL contains:

Flumetasone 0.5 mg

Vehicle cbp1 mL

PRESENTATIONS 50 mL



INDICATIONS

Bovine ketosis. Inflammatory processes associated or not to chemotherapeutic and antimicrobial. Conditions musculoskeletal injuries due to inflammations, when there are no structural changes irreversible Allergic states; tendosynovitis, bursitis, carpalis, myositis, arthritis; eczema. As coadjuvant in infectious processes such as tracheitis, bronchitis, sinusitis, pneumonia and hepatitis; Urogenital infections such as cystitis, nephritis, metritis.

INDICATED SPECIES

Cattle, horses, sheep, goats, dogs and cats

DOSE

Bovine and equine: From 2 to 10 mL.

Sheep and goats: From 1 to 2 mL.

Swine: From 0.2 to 5 mL.

Dogs and cats: 0.1 to 0.5 mL.

The proposed doses can be repeated 12 or 24 hours later, in the opinion of the Veterinarian

ROUTE OF ADMINISTRATION

Intramuscular, intravenous, intraarticular or subcutaneous.

WARNINGS

Do not use this product 15 days before slaughter of animals intended for consumption human. Store in a cool, dry place protected from sunlight. Keep out of reach of children.

CONSULT THE VETERINARY PHYSICIAN YOUR SALE REQUIRES MEDICAL RECIPE