

"AGRICULTURAL USE"



genoxi[®]

BACTERICIDA AGRÍCOLA

Copper Sulfate Pentahydrate + Gentamicin Sulfate + Oxytetracycline Hydrochloride
bactericide / wettable powder

Percentage Composition	% in weigh
Active ingredients:	
Copper Sulfate Pentahydrate	80.00
With a content of 25% copper as an element (Equivalent to 200 g of ai / kg)	
Gentamicin Sulfate:	2.00
Sulfate 0 - 3 - Deoxy - 4 - C - methyl - 3 - (methylamino) - β - L - arabinopyranosyl - (1 → 6) - 0 - [2,6-diamino-2,3,4,6-tetradeoxy - α - D - erythro-hexopyranosyl- (1 → 4)] - 2-deoxy-D-streptamine. (Equivalent to 20 g of ai / kg)	
Oxytetracycline Hydrochloride:	6.00
[4S - (4α, 4aα, 5α, 5aα, 6β, 12a α) monochlorohydrate]	
- 4- (dimethylamino) -1,4,4a, 5,5a, 6,11,12a - octahydro - 3,5,6,10,12,12a, hexahydroxy -6-methyl-1,11-dioxo-2-naphthalenecarboxamide (Equivalent to 60 g ai / kg)	
Inert ingredients	
Antioxidant, anti-caking and vehicle.....	12.00
TOTAL:	100.00
Reg: RSCO-MEZC-1302C-301-002-088	

May be harmful by contact with skin
May be harmful if inhaled

REGISTER HOLDER, FORMULATED AND DISTRIBUTED BY:

LAPISA, SA DE CV
Carr. La Piedad-Guadalajara Km. 5.5
Col. Camelinas, CP 59375,
La Piedad, Michoacán, Mexico.
Tel.: 01 (352) 526 13 00.
www.lapisa.com
MADE IN MEXICO

" ESTE MATERIAL NO ES UNA ETIQUETA, LA FUNCION DE ESTE DOCUMENTO ES DE CARACTER ILUSTRATIVO."



CAUTION

"STOP, READ THE LABEL BEFORE USING THE PRODUCT"

PRECAUTIONS AND WARNINGS OF USE

During the use and handling of this product always use appropriate personal protective equipment: overalls or sleeve shirt long (cotton) and long pants, gloves (neoprene or nitrile), mask, safety glasses, waterproof cap and shoes closed. Wash it with clean water and soap before using your personal protection equipment again. Dry in the sun and store it separately to the clothes of common use and under lock in a cool place

PRECAUTIONS DURING PRODUCT HANDLING

"DO NOT EAT, DRINK OR SMOKE WHEN YOU ARE USING THIS PRODUCT"

"DO NOT TRANSPORT OR STORE NEXT TO FOOD PRODUCTS, CLOTHING OR FORAGE"

"KEEP OUT OF REACH OF CHILDREN, PREGNANT, LACTATION AND ANIMAL WOMEN

DOMESTICOS "

"KEEP IN YOUR CLOSED ORIGINAL CONTAINER AND WITH THE LABEL WHILE NOT USED, REMOVED FROM HEAT AND DIRECT FIRE. KEEP UNDER KEY, IN THE SHADOW, IN A FRESH AND VENTILATED PLACE "

"WASH YOUR HANDS AFTER USING THE PRODUCT AND BEFORE CONSUMING FOOD"

"DO NOT STORE IN HOUSES ROOM"

"DO NOT USE THE PRODUCT WITHOUT THE PROPER PROTECTION EQUIPMENT"

"PERSONS UNDER 18 YEARS OLD SHOULD NOT HANDLE THIS PRODUCT"

"AT THE END OF THE WORK DAY BATH AND PUT CLEAN CLOTHES"

"DO NOT INGEST, INHALE AND AVOID CONTACT WITH EYES AND SKIN"

"DO NOT MIX OR PREPARE THE PRODUCT WITH YOUR HANDS, USE GLOVES, USE A PIECE OF WOOD OR OTHER APPROPRIATE MATERIAL "

"DO NOT UNLOCK THE NOZZLE WITH THE MOUTH"

"MIX THE PRODUCT IN A VENTILATED PLACE

"IN CASE OF INTOXICATION, TAKE THE PATIENT WITH THE DOCTOR AND SHOW THEM THIS LABEL"

FIRST AID

In case of poisoning get medical attention immediately or call the Poison Center (ATOX) for medical assistance 24 hours a day at 01-800-000-2869, meanwhile, it is recommended to apply the following measures of first aid:

Inhalation: bring the intoxicated to a well-ventilated area, with fresh air. If the intoxicated person does not breathe, give breathing

mouth to mouth putting a clean cloth in between. Give one breath every 4 seconds (15 breaths per minute).

Ingestion: if the intoxicant is still conscious, provide one or two glasses of water and induce vomiting by gently touching the bottom of the throat with a finger. In case the patient is unconscious do not administer anything by mouth.

Eyes: wash immediately with plenty of clean water for at least 15 minutes.

Skin: if the person is unconscious, using gloves immediately remove contaminated clothing and wash the skin with plenty water and soap. If you are aware, immediately bathe.

RECOMMENDATIONS TO THE DOCTOR

This product is a bactericide, a mixture of Systemic Aminoglycosides and Tetracyclines with Copper Sulphate.

Signs and symptoms of poisoning: Upset stomach and diarrhea. Product slightly irritating on the skin.

Antidote: No specific antidote is known. Use symptomatic treatment

MEASURES OF PROTECTION TO THE ENVIRONMENT


"DURING THE HANDLING OF THE PRODUCT, AVOID THE CONTAMINATION OF SOILS, RIVERS, LAGOONS, STREAMS, DAMS, CHANNELS OR WATER TANKS, NOT WASHING OR SEWING IN THE PESTICIDE RESIDUES OR EMPTY CONTAINERS".

"MANAGE THE EMPTY CONTAINER AND THE RESIDUES OF THE PRODUCT AS PROVIDED IN THE GENERAL LAW FOR THE PREVENTION AND INTEGRAL MANAGEMENT OF WASTE, ITS REGULATIONS OR THE MANAGEMENT PLAN FOR

EMPTY PESTICIDE PACKAGING, REGISTERED BEFORE THE SEMARNAT "

"IN CASE OF ACCIDENTAL FALL OF THE PRODUCT, PERSONAL PROTECTIVE EQUIPMENT SHOULD BE USED AND RECOVER THE PRODUCT THAT IS USEFUL UNTIL WHERE POSSIBLE, AND OTHERWISE COLLECT IT IN A HERMETIC CONTAINER AND TAKE IT TO THE AUTHORIZED DANGEROUS WASTE DISPOSAL CENTER MORE NEAR".

"PERFORM THE TRIPLE WASHING OF THE EMPTY CONTAINER AND VIEW THE RINSE WATER IN THE DEPOSIT OR CONTAINER WHERE PREPARE THE MIX FOR APPLICATION "



"USE EXCLUSIVELY IN THE CULTIVATION AND DISEASE HERE RECOMMENDED"

INSTRUCTIONS FOR USE

"ALWAYS CALIBER YOUR APPLICATION TEAM"

CROP	PLAGUE	DOSAGE kg / ha
Agave (*)	Cause of Rotting of the heart (<i>Pantoea agglomerans</i>) (<i>Erwinia herbicola</i>)	11.5 - 2.0 kg Application to pens and bases of the same.
Eggplant Chile Pepper Jitomate Potato	<i>Bacterial stain</i> (<i>Xanthomonas vesicatoria</i>) <i>Bacterial wilt</i> (<i>Ralstonia solanacearum</i>)	0.2 - 0.4 kg / ha Perform 2 to 3 applications at foliage with 7 days interval when the first ones are detected symptoms of the disease in a volume of 500 to 600 L of water per ha.
Cucumber Melon Watermelon Zucchini Pumpkin Squash	Angular spot (<i>Pseudomonas syringae</i> <i>Pv lachrymans</i>)	0.3- 0.5 kg / ha Perform 2 to 3 applications at foliage with an interval of 5 to 7 days depending on the pressure of the disease Y from the environmental conditions in a volume of 460 to 550 L of water per ha.

(*) There is no safety interval, so exclusive use in plantations under 5 years is recommended.

TIME OF REENTATION TO THE TREATED ZONES

DO NOT ENTER THE TREATED AREA BEFORE 24 HRS

METHOD FOR PREPARING AND APPLYING THE PRODUCT

Open the container carefully at the top and measure the amount of product indicated on this label, mix in the spray tank with half the recommended water, then add the other half of water. Keep the mixture in agitation.

After using your application equipment, wash it with soap and water. Let the water circulate throughout the equipment, in order to eliminate as much as possible waste. Do not open the mouthpiece with your mouth.

CONTRAINDICATIONS

It is recommended to be used exclusively in plantations under 5 years old.

PHYTO-TOXICITY

This product is not phytotoxic to the crop and in the doses indicated here, if it is applied according to the recommendations of this label.

INCOMPATIBILITY

This product should not be mixed with other products.

RESISTANCE MANAGEMENT

"TO PREVENT THE DEVELOPMENT OF RESISTANT POPULATIONS, ALWAYS RESPECT THE DOSE AND FREQUENCIES OF APPLICATION; AVOID REPEATED USE OF THIS PRODUCT, ALTERNATING IT WITH OTHER CHEMICAL GROUPS OF DIFFERENT MODES OF ACTION AND DIFFERENT MECHANISMS OF DETOXIFICATION AND THROUGH THE SUPPORT OF OTHERS CONTROL METHODS".

WARRANTY

LAPISA, SA de CV guarantees that said product will be composed of the specified ingredients. Does not accept responsibility some for damages to persons or damages to the property of others that could result from its use if it is not fulfilled according to the instructions of the label

