

Lapicor®

Dexamethasone in Solution for injection

REG. SAGARPA Q-2083-010

FORMULA

Each mL contains:

Dexamethasone base 2 mg (As 21 Dexamethasone phosphate)

Vehicle cbp 1 mL

PRESENTATIONS

Ampules with 20 mL and 50 mL.



DESCRIPTION

Dexamethasone is an anti-inflammatory agent, gluconeogenic and anti-stress, synthetic analogue of prednisolone.

INDICATIONS

Due to its potent gluconeogenic effect, it is indicated in the treatment of primary bovine ketosis; for his powerful anti-inflammatory effect is indicated in the treatment of acute or chronic inflammations in any animal species: various arthritis and osteoarthritis, rheumatism, myositis, dermatitis, excemas, pruritus etc. As well It is indicated as supportive therapy in infections such as boarding fever, mastitis, metritis and pyelonephritis, combined with a suitable primary antibiotic therapy. Also as an anti-inflammatory in cases of acute mastitis, pneumonia, laminitis and retention of placenta

USE

All domestic mammals.

DOSAGE, MODE AND ROUTE OF ADMINISTRATION

Bovine: 5 to 20 mg (2.5 to 10 mL) intravenously or intramuscular for 3-5 days.

Equines: 5 to 10 mg (2.5 to 5 mL) intravenously, intramuscularly or intra-articular for 3-5 days.

Porcine and Ovi-Caprines: from 5 to 10 mg (2.5 to 5 mL) intravenously or intramuscularly for 3-5 days.

Canines and Felines: 0.25 to 1 mg (0.125 to 0.5 mL) intravenously, intramuscularly or intra-articularly for 3-5 days.

WARNINGS

The use of any corticosteroid at the end of pregnancy (last third) can accelerate the calving process in cattle, sheep and goats.

This product should not be used 5 days before slaughter of animals destined for human consumption. Milk from treated cows should not be used until 72 hours after treatment.

CONSULT A VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION