

Lapigen 50

Gentamicin in Injectable Solution

REG. SAGARPA Q-2083-042

FORMULA

Each mL contains:
Gentamicin 50 mg base
(as Gentamicin sulfate)
Vehicle c.b.p. 1 mL

PRESENTATIONS

Bottle with 250 mL.



INDICATIONS

Broad-spectrum antibiotic indicated in the treatment of diseases caused by Gram-positive and Gram-negative germs in cattle, swine, horses, dogs and cats. For the treatment of bacterial infections of the respiratory, gastrointestinal and genitourinary tracts as well as mastitis.

DOSAGE AND ADMINISTRATION

Respiratory infections and septicemia: 5 mg / kg / day intramuscular 3 to 5 days.
Intestinal and genitourinary infections: 4 mg / kg / day intramuscularly for 3 to 5 days.
Cervicitis, metritis, endometritis in cows: from 200 mg to 1,000 mg diluted in physiological saline. Intrauterine application.

Swine: for the prevention and treatment of colibacillosis and intestinal infections, inject 5 mg per piglet intramuscularly as a single dose for 3 consecutive days.

Poultry: for the prevention of early mortality in one-day-old chickens caused by E. coli, E. aerógenes, Salmonella and P. aeruginosa.

Mares with metritis, endometritis and cervicitis: use intrauterine from 500 mg to 2500 mg daily from 3 to 5 days during the heat season when the cervix is relaxed.

WARNINGS

Do not use this product 60 days before the slaughter of animals intended for human consumption.

Do not use milk from animals treated with this product for human consumption, until 4 days after the last application.

Do not use in pregnant females.

Do not apply in cases of renal failure or after prolonged treatment with other aminoglycosides.

Do not use together with diuretics.

Store in a cool, dry place, protected from light.

CONSULT THE VETERINARY PHYSICIAN
YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION