

LAPIMICINA®

Injectable oxytetracycline

REG. SAGARPA Q-2083-004

FORMULA

Each mL contains:

Oxitetraciclina base 50 mg (as oxytetracycline hydrochloride)

Vehicle cbp 1 mL



INDICATIONS, INDICATED SPECIES AND SUSCEPTIBLE ORGANISMS

Broad-spectrum antibiotic indicated in all species for the treatment of pneumonic problems, enteritis, leptospirosis, as well as various septicemic infectious processes that are caused by germs susceptible to oxytetracycline.

SPECTRUM OF ACTION

Pasteurella spp., Haemophilus spp., Moraxella bovis, Fusobacterium necrophorum, Escherichia coli, Actinobacillus lignieresii, Leptospira spp., Staphylococcus spp., Streptococcus spp., Etc.

DOSAGE AND ROUTE OF ADMINISTRATION

Administer at a rate of 5 to 10 mg per kg of live weight for 4 consecutive days. It can be administered by any means as necessary.

INCOMPATIBILITY

Because bacteriostatic drugs interfere with the bactericidal action of penicillin, do not administer oxytetracycline concurrently with penicillin.

PRECAUTIONS

The use of the product may cause transient local irritation manifested by inflammation and discoloration at the injection site. The pain produced by the injection can be minimized by heating the solution at body temperature before administration and avoiding a bad injection technique. After treatment, animals may exhibit transient hemoglobinuria. If an improvement in clinical signs is not observed in 24-48 hours, consult your veterinarian. Do not use the product for more than 4 consecutive days. Exceeding dosage levels or duration of treatment, will cause tissue waste is not eliminated in the period of suspension. Do not use in growing animals or during pregnancy. Do not administer together with diuretics or intravenous calcium. Store in a cool, dry place protected from sunlight. Keep out of the reach of children.

RETIREMENT PERIOD

This product should not be used 30 days before the slaughter of animals destined for human consumption. Do not use milk from cows treated for up to 72 hours for human consumption after the last application.

PRESENTATIONS

Bottles with 100 and 500 mL.

CONSULT THE VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION