

LAPI-RESP FT®

Antibiotic premix
REG. SAGARPA Q-2083-161

FORMULA

Each kg contains:
Tylosin tartrate 110 g
Florfenicol 40 g
Excipient cbp 1 kg

PRESENTATIONS

20 Kg.

USE

Poultry and Pigs

INDICATIONS

Antibiotic premix indicated for the prevention and treatment of respiratory diseases and digestive pigs of all ages and in fattening birds, caused by Salmonella spp, Escherichia coli, Pasteurella haemolytica, Pasteurella multocida, Haemophilus somnus, Haemophilus parasuis, Streptococcus suis, Mycoplasma spp and Actinobacillus pleuropneumoniae.

DOSAGE AND ADMINISTRATION

Species	Preventive treatment	Therapeutic Treatment
Birds	Administer 500 g of Lapi- Resp FT by tonne of finished food, (20 ppm of florfenicol and 55 ppm tylosin), during a period of 5 to 7 days	Administer 1 to 2 Kg of Lapi- Resp FT by ton of finished food, (40 to 80 ppm of florfenicol and 110 to 220 ppm of tylosin), for a period of 5 to 7 days.
Pigs	Administer 500 g of Lapi- Resp FT by tonne of finished food, (20 ppm of florfenicol and 55 ppm tylosin), during a period of 5 to 7 days.	Administer 1 Kg of Lapi- Resp FT by tonne of finished food, (40 ppm of florfenicol and 110 ppm tylosin), during a period of 5 to 7 days.

ROUTE OF ADMINISTRATION

Oral, mixed with the finished food.

WARNINGS

Do not use this product in laying birds.
You should not use this product 7 days before the slaughter of birds and pigs intended for human consumption.
Keep in a cool and dry place.
Keep out of reach of children.

CONSULT A VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION

