

Laptil-300

Tilmicosin in injectable solution

REG. SAGARPA Q-2083-169



FORMULA

Each mL contains:

Tilmicosin (As phosphate) 300 mg

Vehicle, cbp 1 mL

PRESENTATIONS

500 mL bottle.

INDICATIONS

Laptil-300 is indicated in the treatment of diseases respiratory mainly associated with Mannheimia (Pasteurella) haemolytica, Pasteurella multocida and other organisms susceptible to tilmicosin

USE

Sheep and cattle

DOSE

In adult bovines, heifers, calves: Administer a single dose of 10 mg / kg of weight (1 mL / 30 kg of weight).

In sheep over 15 kg, administer a dose of 10 mg / kg of weight (1 mL / 30 kg of weight).

Do not administer more than 10 mL per injection site.

ROUTE OF ADMINISTRATION

Apply only subcutaneously, preferably apply to the neck table.

WARNINGS

You should not use this product 42 days before slaughter of cattle and sheep intended for human consumption.

Do not use in lactating cattle.

Do not apply with automatic injection equipment. Avoid contact with skin and mucous membranes through the use of adequate protection (gloves, glasses).

Do not administer intravenously since it can cause the death of the animal.

Do not administer intramuscularly because it can cause local reaction and represent losses by cuts to the channel.

The injection of tilmicosin in humans, pigs, horses and goats can cause death.

Toxicity in man does not occur by manipulation of the product, only by injection or ingestion accidental. In this case, go immediately to the doctor.

The cardiovascular system is the most affected by the toxicity, which could be due to the blockage of the channels of calcium.

The use of epinephrine as a therapy is contraindicated.

Store in a cool, dry place protected from light.

Keep out of reach of children.

CONSULT A VETERINARY

VETERINARY USE