

Larvakill Premix

Larvicide premix for food

RSCO-PEC-INAC-0173-310-001-001



PRESENTATION

20 kilos sack

DESCRIPTION

It is a synthetic growth regulator of insects with activity against flies and fleas. Its manifest activity by contact, interfering in the process of moulting in the larval stages that prevents normal pupation, causing pupae deformed that prevents the flies from hatching and adult fleas.

Prevents larval development and pupation of the Stable fly *Stomoxys calcitrans*, fly of the Horn *Haematobia irritans*, small fly domestic *Fannia canicularis*, manure fly of chickens *Fannia pusio*, housefly *Musca domestica*, fruit fly *Drosophila melanogaster* and other dipterous larvae that reproduce in manure or waste in fermentation, as well as in flea larvae like *Echidnophaga gallinacea* and *Pulex irritans* that they develop in the hens' bed commercial and reproductive layers

COMPOSITION

Active ingredient % in weigh

N- cyclopropyl-1,3,5-triazine-2,4,6-triamine

Containing not less than: 1 (equivalent to 10 g ai / kg)

Inert ingredients:

Vehicles and stabilizers

Containing no more than: 99

TOTAL 100

SPECTRUM OF ACTION

Fly larvae

Stable fly *Stomoxys calcitrans*

Horn fly *Haematobia irritans*

Small housefly *Fannia canicularis*

Chick manure fly *Fannia pusio*

Housefly *Musca domestica*

Fruit fly *Drosophila melanogaster*

Flea larvae

Flea of the hen - *Echidnophaga gallinacea*

Man's flea - *Pulex irritans*

HOW TO USE

Larvakill Premix is added to balanced feed of the layers during its elaboration. The hens ingest it in each daily ration, being eliminated with the feces, where it acts on the larvae of the flies and fleas that they develop in the chicken manure and in the bed of the birds respectively.

DOSE

Mix 500 grams of Larvakill Premix by ton of balanced feed, when starting the fly season for 4 to 6 weeks followed and suspend for the same period (4 to 6 weeks) and so on.

The larvicidal action of Larvakill Premix begins to be observed three to five days after the application.

ADVANTAGE

Different mechanism of action for being a regulator of the growth of insects that acts inhibiting the moult and the pupation.

Used within a comprehensive program of adulticides and larvicides prevents reinfestation of flies and fleas.

Used within a comprehensive program of adulticides and larvicides prevents reinfestation of flies and fleas

PRECAUTIONS

Withdrawal time: Do not consume poultry meat treated up to three days after finishing the addition of it.

Avoid contaminating water sources with residues of washing or cleaning.

Keep out of reach of children and animals.

Store in a cool and dry place.

Avoid the entry of water to the maximum manure and the bed to not dilute the Larvakill Premix below the minimum concentration of larvicidal effectiveness.

CONSULT THE VETERINARY PHYSICIAN