

LOMO PON S[®]

Insecticide, mosquicida

Reg. RSCO-PEC-INAC-0156-395-378-007



FORMULA

ACTIVE INGREDIENT % in weigh

PERMETHRINE: 3-phenoxybenzyl (1RS) -CIS, trans -3- (2,2-dichlorovinyl)
-2,2-dimethyl cyclopropane carboxylate

(Equivalent to 64.9 gr of AI / L) not less than: 7.40%

BUTOXIDE OF PIPERONILO no less than: 7.40%

AROMATIC INGREDIENTS VEHICLE no more than: 85.20%

TOTAL 100.00%

PRESENTATIONS

1 and 5 liters.

INDICATIONS

It is a product based on a pyrethroid and a synergist, with action on populations of pallet flies or the horn. Permethrin is a pyrethroid that has neurotoxic action on the nerve membranes of ectoparasites, causing excitement, coordination disorders, paralysis and death. The synergistic effect of piperonyl butoxide on the resistance enzymes prevents the degradation of the permethrin multiplying its action

USE LOMO PON S[®] is a product formulated for pour on application. It has action on the main Cattle parasites: Horn or palette fly : *Haematobia irritans* . Stable fly or brave fly: *Stomoxys calci-* trans House fly, horseflies and mosquitoes. Sucking lice : *Haematopinus eurysternus*, *H. suis*, *Lignogatus vitulli*, *Solenopotes capillatus* . Chewing lice: *Melophagus* spp; *Damalinia* spp .

DOSAGE AND ROUTE OF ADMINISTRATION

Apply in a strip from the cross to the rump. In animals weighing less than 200 kg, apply 10 mL. In animals weighing more than 200 kg, apply 20 mL..

WARNINGS

Period of withdrawal in meat and milk: zero days. Store in a cool, dry place. Keep out of reach of children.

CONSULT A VETERINARY
VETERINARY USE