

Mamitrol Fuerte

Intramammary Antibiotic Infusion

REG. SAGARPA Q-2083-112

FORMULA

Each 10 mL syringe contains:

Dihydrostreptomycin (as sulfate) 0.25 g

Neomycin (as sulfate) 0.1 g

Polymyxin (as sulfate) 80,000 IU

Flumetasone 0.25 mg

Vehicle cbp 10.0 mL



PRESENTATIONS

Box of 12 syringes with 10 mL each, accompanied by cannula specially designed for the udder.

INDICATIONS

The product has been designed for the treatment and control of sub-acute, acute mastitis or Chronic diseases, accompanied by pain and inflammation, in cattle, originated by microorganisms susceptible to Dihydrostreptomycin and Neomycin. The potent corticosteroid facilitates the diffusion of antibiotics in inflamed tissues.

The medication is done intramammary after a good milking of the affected room, applying the content of a 10 ml syringe, with a light massage upwards.

INDICATED SPECIES

Bovines, goats.

DOSE

Cattle: one syringe in each infected room.

The treatment can be repeated every 12 hours, for 3 or 4 days according to the evolution of the suffering

ROUTE OF ADMINISTRATION

Administer intramammary after milking, apply in the infected room with a gentle massage

WARNINGS

Milk from animals treated with this product should not be used for consumption until 4 days after the last application.

Store in a cool, dry place protected from sunlight.

Keep out of reach of children

CONSULT A VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION