

Mastrix

Intramamaria Solution

REG. SAGARPA Q-2083-076

FORMULA

Each ml contains:
Lincomycin 20.0 mg
Neomycin 20.0 mg.
Dexamethasone 0.1 mg.
Vehicle cbp1.0 mL



INDICATIONS

Mastrix is an aqueous solution for intramammary infusion based on 20% Lincomycin, 20% Neomycin and Dexamethasone, for the treatment of subacute, acute or chronic mastitis in bovines produced by microorganisms susceptible to the formula.

DESCRIPTION

The product has been designed for the treatment and control of sub-acute, acute mastitis or Chronic diseases, accompanied by pain and inflammation, in cattle, originated by microorganisms susceptible to Lincomycin and Neomycin. The potent corticosteroid facilitates the diffusion of antibiotics in inflamed tissues .

The medication is done intramammary after a good milking of the affected room, applying the content of a 10 mL syringe, with a light massage upwards.

The treatment can be repeated every 12 hours, for 3 or 4 days according to the evolution of the suffering Lincomycin being a broad spectrum antibiotic with action against gram positive and gram germs negative, its association with Neomycin, also broad-spectrum create a wide range of antibacterial action against the main germs involved in mastitic processes, such as

Gram positives:

Staphylococcus aureus.
Streptococcus agalactiae.
Streptococcus dysgalactiae.
Streptococcus uberis
Streptococcus pyogenes.

Gram negative:

Escherichia coli
Klebsiella spp.
Shigella spp.
Pseudomonas spp.
Salmonella spp.
Mycoplasma spp.

During the process of this disease the inflammation of the gland accompanied by edema occurs, heat and pain. Dexamethasone helps to alleviate these signs, in addition to allowing a better diffusion of the antibiotics through inflamed tissues, decreasing the recovery time.

INDICATED SPECIES

Bovine.

DOSE

The medication is done intramammary after a good milking of the affected room, applying the content of a 10 mL syringe, with a light massage upwards. The treatment can be repeated on every 12 hours, for 3 or 4 days according to the evolution of the condition

ROUTE OF ADMINISTRATION

Intramamaria

WARNING

Milk from animals treated with this product should not be used for up to 4 days after the last application.

Store in a cool, dry place protected from sunlight.

Keep out of reach of children.

PRESENTATIONS

Box with 24 syringes of 10 mL

CONSULT A VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION