

# MAXI-LAP®

MAXI-LAP® is a premix based on an  $\beta$ -agonist (Zilpaterol) at a concentration of 4.8%, which acts as a promoter of growth and performance for finish fattening cattle.

REG. SAGARPA Q-2083-076

## FORMULA

Each Kg Contains:  
Zilpaterol Hydrochloride 48 g  
Excipient c.b.p. 1 kg



## INDICATIONS

MAXI-LAP® is indicated for the improvement of the degree of performance of the carcass, decreasing the deposit of fat, as it improves the gain of weight and the alimentary conversion in bovine of fattening.

## SPECIES

Bovine fattening.

## DOSE

The dose of zilpaterol hydrochloride is 0.15 mg / kg of live weight per day, which corresponds to a food concentration of 6 ppm for an animal with an average weight of 400 kg that ingests approx. 10 kg of finished food (with a content of 90% dry matter) per day during the last 30 days of the fattening in corral. Therefore, in practice, it is recommended finished food with a content of 90% dry matter.

## ROUTE OF ADMINISTRATION

Oral mixed in the food.

## WARNINGS

Dosing should be done exactly to respect the recommended dose.

The product should be stored in a cool, dry place at room temperature (<25 ° C).

For the handling of MAXI-LAP® follow the recommendations for the preparation of premix: use a mask with filter, gloves, protective glasses and overalls.

Avoid direct inhalation of the product and direct contact with the skin and eyes.

Store away from food and keep out of reach of children.

Do not smoke or consume food during handling.

In case of contact with the skin, rinse with plenty of running water.

In case of accident, call the doctor.

In case of accidental ingestion, a B2 blocker such as propranolol can be administered under strict medical supervision.

## WITHDRAWAL PERIOD

This product should not be used 72 hours before the slaughter of animals destined for human consumption.

CONSULT A VETERINARY

REQUIRE A MEDICAL RECEIPT

EXCLUSIVE USE OF THE VETERINARY