

NEUMO 200®

Oxytetracycline premix

REG. SAGARPA Q-2083-044

FORMULA

Each kg contains 200 g

Oxytetracycline Excipient c.b.p.1000 g



PRESENTATIONS

Sack with 25 Kg.

INDICATIONS AND DOSES

For the prevention and treatment of diseases caused by germs sensitive to the action of oxytetracycline. Mycoplasma hyopneumoniae, Clostridium spp., Haemophilus spp., Pasteurella multocida, Leptospira spp., Erysipelothrix rhusiopathie, among others.

SWINE

The premix acts as a growth promoter when administered continuously. Also in the prevention of respiratory diseases and as a treatment for leptospirosis, mycoplasmosis and bacterial respiratory problems.

Generally Neumo 200® is strategically combined with other antibiotics to expand the spectrum of action and the effectiveness of the treatment.

POULTRY

For the prevention of chronic respiratory disease, infectious synovitis, avian cholera, mycoplasmosis.

Program	Preventive Kg/Ton	Active Grams	7 i fUlj Yg Kg/Ton	Active Grams
Øæ^} ã * Á&@A ^}	0.5 - 1.0 Kg Ø[{ Á Á Á Á Áæ • D	100-200 g	1.5 – 2.5 Kg (From 10 to 15 days)	300 - 500 g
Ú[• ç !^Áãã•	0.5 - 1.0 Kg Ø[{ Á Á Á Á Áæ • D	100-200 g	1.5 – 2.5 Kg (From 10 to 15 days)	300 - 500 g
Øæ^} ã * Áã•	1.5 Kg Ø From ì Á Á Á Á Áæ • D	300 g	2.0 – 3.0 Kg (From 8 to 10 days)	400 - 600 g
Ói^ãã * Á[, •	1.5 Kg Ø[{ Á Á Á Á Áæ • Á[• ð { [} D	300 g	2.0 – 2.5 Kg (From 8 to 10 days or 15 - 5 * - 15 days)	400 -500 g

*Days without medication

CATTLE

As a promoter of growth, prevention and treatment of bacterial pneumonias, boarding fever, hemorrhagic septicemia, bacterial diarrhea, pododermatitis.

In general, it is dosed:

Preventive: 5 g. per head per day for a period between 7-10 days.

Healing: 10 g. per head per day for 7 days or more if necessary.

In case of giving concentrate or a rationed diet per day, calculate the consumption per day of the animal, calculate how many heads are fed with one ton, multiply the number of heads per grams of active ingredient and multiply the result by 5 to obtain how much finished product should be added per ton of food.

ROUTE OF ADMINISTRATION:

Oral mixed with food.

WARNINGS:

Do not use this product 4 days before the slaughter of animals intended for human consumption. Keep out of reach of children.

Store in a cool, dry place, protected from light.

PRESENTATIONS:

Sack with 25 Kg.

CONSULT THE VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION