

Oviver®

Endectocide and Fasciolicide of broad spectrum in solution injectable

REG. SAGARPA Q-2083-157



FORMULA

Each mL contains: ivermectin 5.0 mg

Closantel 125.0 mg

PRESENTATION

Plastic bottles of 100 mL.

Vehicle, cbp 1.00 mL

Therapeutic indications:

For control parasitic from nematodes gastroenteric, pulmonary, fasciola, larvae flies such as Oestrus ovis, Hipoderma and external parasites such as sucking lice, ticks and Mites.

DOSE

Sheep and goats 1 mL per 25 kg of live weight

ROUTE OF ADMINISTRATION

Subcutaneous, in the table of the neck or in the armpit. OVIVER® can be administered with a syringe of type automatic or conventional.

Management recommendations

Lambs: At month of age, and at weaning.

Bellies, stallions and replacements:

Every 3 or 4 months depending on the signs of the "worm of the nose "or estrosis.

WARNINGS FOR EXCLUSIVE USE ON SHEEP AND GOATS.

This product must not be used 28 days before the slaughter of animals destined for consumption human.

Do not administer this product to Caprinos producers of milk that is intended for consumption human.

Some cases of animals with transient signs have been observed after administration subcutaneous, such as: slight inflammation at the site of injection and transient pain in sheep; these reactions disappear without the need for treatment. Aquatic organisms are highly sensitive to the toxicity of Ivermectin, so the recipients Several and any residual contents must be disposed of safely by burying or incinerating them.

It should not be applied intramuscularly or intravenously. Keep out of reach of children. Store in a cool, dry place protected from light.

CONSULT A VETERINARY.

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION.