

# Influenza Porcimune®

Oily suspension for injection.

**REG. SAGARPA B-2083-025**

## FORMULA

Crops swine flu H1N1 and H3N2 inactivated serotypes and adyuvantados.

## PRESENTATION

50 dose /100 mL.



## DESCRIPTION

Swine Influenza is a highly contagious disease of pigs, is associated as the primary agent of porcine respiratory disease complex. The virus is cosmopolitan distribution, outbreaks of the disease are characterized by high morbidity and mortality although usually less than 5%, is a primary agent allowing entry to other pathogens into the body. The main symptoms of influenza in pigs include sudden onset of fever, depression, coughing, runny nose and eyes, sneezing, difficulty breathing, swelling of the eyelids and loss of appetite. Vaccination is one of the tools to control the disease in swine herds. In Mexico, the H1N1 and H3N2 serotypes are primarily those who have been associated boxes influenza in pig farms.

## INDICATIONS

Prevention and control in healthy pigs from disease associated with swine flu virus H1N1 and H3N2 serotypes.

## DOSE AND ADMINISTRATION MODE

2 mL intramuscularly administered to healthy pigs from the fourth week of age. Revaccination with a dose to the second or third week after the initial vaccination.

## WARNINGS

Shake the bottle before use. Once broached use the full content.

Keep refrigerated at a temperature between 2 and 7 C, do not freeze. Any biological can cause anaphylactic reactions. In this case the use of epinephrine followed by appropriate supportive therapy is recommended. Do not vaccinate 21 days before slaughter of animals intended for human consumption. Discard the container safely bury yourself or incinerate. Keep out of reach of children

CONSULT THE VETERINARY PHYSICIAN