

Porcimune® SOA

Inactivated vaccine for the prevention and control of Blue Eye Disease in healthy swines.

REG. SAGARPA B-2083-017

FORMULA

Each mL contains:

Porcine rubulavirus strain LPMV-SOA-CI and LPMV-SOA-CIII minimum 1x10⁶ U.F.P. Chemically inactivated and emulsified with an oil adjuvant.



INDICATIONS

Inactivated vaccine for the prevention and control of Blue Eye Disease in healthy swines.

INDICATED SPECIES

Healthy swine.

DOSE

The dose is 2 ml and must be applied under the following schedule:

Piglets: Apply a dose between 6 and 8 weeks of life and repeat the dose two weeks later. Breeding Foot

Replacements: Two applications during adaptation or quarantine. (With interval of 2 weeks). Females:

Females can be vaccinated safely at any time.

The following vaccination programs are suggested:

Initially two applications with an interval of 2 weeks and then every 4 to 6 months. (bed sheet). Initially

two applications with an interval of 2 weeks and then an application 3 weeks before delivery. Stallions:

Initially two applications with an interval of 2 weeks and then every 4 to 6 months.

ADMINISTRATION MODE

Apply by deep intramuscular injection.

WARNINGS

Once the container is opened, use the contents completely. Discard the excess product.

Shake the bottle before use.

Store in refrigeration at a temperature between 2° and 7°C.

Any biologic can cause anaphylactic reactions. In such a case, administer antihistamines.

Do not vaccinate 21 days before the slaughter of animals intended for human consumption

Do not vaccinate 21 days before the sacrifice.

FOR EXCLUSIVE USE IN THE STATES AUTHORIZED BY SAGARPA.