

PROVIDEAN PRIMOTEC AT

High titre vaccine against Canine Parvovirus.

REG. SAGARPA B-2083-028

FORMULA

Each dose of 1 mL contains:

Live attenuated virus (50% DICC / dose): Parvovirus ($\geq 10^8$).



PRESENTATIONS

20 bottles with 1 dose each.

INDICATIONS

Vaccination of dogs in good general health is recommended, with proper nutritional development, deworming, without febrile status and out of stress due to adverse environmental conditions.

Primary vaccination: In puppies from 6 weeks of age apply a three-dose vaccination program, maintaining an interval of 3 weeks in each of them. It is advisable to have finished the vaccination program before 12 weeks of age to induce the window period of exposure of the puppies during their socialization phase.

Revaccination: Apply a booster dose every year

DOSAGE AND ADMINISTRATION

Canines: 1 mL.

Shake the vial before use. Using sterile needle and syringe aseptically extract the contents and administer 1 mL intramuscularly or subcutaneously.

WARNINGS

Keep out of the reach of children. Keep the product in its original container between 2° and 8° C. Do not freeze. Use all the content Use sterile disposable syringes for their application.

Avoid vaccination of pregnant females. All vaccines can cause anaphylactic reactions, in which case apply corticosteroid.

VETERINARY USE

CONSULT THE VETERINARY DOCTOR

Made in Argentina by:

Tecnovax S.A, Luis Viale 2835 (C1416DNU) Ciudad Autónoma de Bs. As, Argentina.

SENASA: Certificado N° 06-147. Est. Elaborador N° 8520.

Tel./Fax: (+54) (11) 4585-9695 /4899-0568.

www.tecnovax.com

D.T (Director Científico): Dra. Marcela Iglesias M.P 7140.