

SECACEF®

Intramammary benzathine cephapirin

REG. SAGARPA Q-2083-088

FORMULA

Each syringe with 10 mL contains:

Cefapirin 300 mg (as Benzathine cephapirin)

Vehicle cbp 10 mL



PRESENTATIONS

Box with 12 syringes of 10 mL

INDICATIONS

Infusion intramammary for prevention and treatment of bovine mastitis in dry period. Benzathine Cefapirin has been shown to have a bactericidal action on susceptible organisms, as they are: Streptococcus agalactiae and Staphylococcus aureus, also against that show resistance to penicillin. Benzathine cephapirin is a cephalosporin that has a wide range of activity against Gram positive and gram negative organisms. This is derived biosynthetically from acid 7- aminocephalosporin.

USO

Bovine

DOSAGE AND ROUTE OF ADMINISTRATION

Use exclusively in dry cows. Apply the contents of one syringe in each room at the end of the last milking of the lactation period. Intramamaria

WARNINGS

Do not use meat from animals treated with this for human consumption

Product up to 5 days after the last application.

Do not consume milk from cows treated 120 hours after delivery.

Store in a fresh and dry place.

Keep out of the reach of children

CONSULT THE VETERINARY

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION

VETERINARY USE