

Sulfatropin® Plus

Soluble Powder

REG. SAGARPA Q-2083-043

FORMULA

Each 100 g contains: Sulfachloropyridazine sodium 10g Trimethoprim 2g
Vehicle c.b.p. 100g



PRESENTATIONS

Buckets of 5 kg.

INDICATIONS

It is indicated in the treatment of the infectious diseases of birds, pigs and calves caused by organisms susceptible to the action of the formula.

Poultry: colibacillosis, ECR, avian typhus, pulorosis, salmonellosis, infectious coryza, avian cholera, etc. Pigs: colibacillosis, colienterotoxemia, bacterial diarrhea, bronchopneumonia, enzootic pneumonia, atrophic rhinitis, MMA complex.

Calves: diarrhea, gastroenteritis, colisepticemia, bronchopneumonia, polyarthritis, diphtheria of calves, infections of the umbilicus.

ROUTE OF ADMINISTRATION

Oral, dissolved in water or mixed with the finished food.

DOSE

In birds and pigs, the following tables serve as a reference.

POULTRY	Age in weeks	Amount of Sulfatropin® Plus soluble	
		Per ton of feed	Per 100 liters of water
For 3 to 6 days			
Broilers	1-4	1.5 kg	100 g
	5-8	2.5 kg	150 g
Broilers	1-8	2.0 kg	100 g
	9-20	3.0 kg	150 g
Laying hen	20	2.5 kg	150 g
Turkeys	20	2.5 kg	100 a 150 g

SWINES	Animal weight	Amount of Sulfatropin® Plus soluble	
		Per ton. of food	Per 100 litres of water
Suckers	Up to 9 kg	7-10 kg	200-350 g
Weaners	10-23 kg	4-5 kg	100-200 g
Fattening pigs	24-105 kg	5-7 kg	200-300 g
Swine before or after delivery MMA	175 kg (average)	11 kg	
Farrowing sows	175 kg (average)	5-7 kg	100-200 g

CALVES:

Administer 10 g of Sulfatropin® plus for every 50 kg of live weight for 5 to 10 days, dissolved in the milk. Divide the dose and administer half in the morning and half in the afternoon.

WARNINGS

Do not use this product 4 days before the slaughter of animals intended for human consumption.

CONSULT THE VETERINARY PHYSICIAN

YOUR PURCHASE REQUIRES A MEDICAL PRESCRIPTION