

TICOFF®

Tickicide and mosquicida for livestock

REG. RACO INAC 111-12-2083-001



FORMULA

Percentage composition% by weight Cypermethrin: 2 cyano-3 Phenoxybenzyl, Trans 3-(2,2-dichlorovinyl -3,2) Propane carboxylate cycle. (Equivalent to 200 g of i.a./L at 20 ° C)

No less than: Inert ingredients: 20.00% Emulsifier, Not more than: 80.00%

solvent TOTAL: 100.00%

PRESENTATIONS

30, 100 mL, 1 L.

PHARMACOLOGICAL PROPERTIES

Cypermethrin is a synthetic pyrethroid used as an insecticide to control infinity of pests, directly on animals or on crops, including cotton moths, fruits, and vegetable crops. It is also used in cracks and crevices for the control of insects in stores, warehouses, industries, houses, apartments, nurseries, laboratories, ships, trains, buses, trucks, and airplanes. It can also be used in schools (except food areas) hospitals, hotels, food processing plants and as a repellent barrier treatment in horses.

When applied to animals, it is absorbed by skin in a low proportion (1% in 24 hours in rats), although its detoxification is very rapid and does not accumulate in animal tissues or in the environment, unlike pesticides. It is rapidly metabolized in the liver by hydrolysis and oxidation reactions

USE RECOMMENDED BY THE MANUFACTURER

Ticoff acts on all the stages of the ticks of one or several hosts with an excellent efficacy and prolonged residual effect (10-14 days). Ticoff also controls other external parasites that affect livestock causing irritation, pain, and discomfort, such as: barn fly, paddle fly, face fly, as well as horseflies, lice, and mites.

DOSAGE AND ADMINISTRATION MODE

Immersion bath:

Load: Cover the volume of the bath, use 1 liter of Ticoff for every 1000 liters of water. Pre-mix the necessary amount of Ticoff in an equal amount of water and add this mixture to the water of the immersion bath.

Recharge: Add 1.5 liters per 1000 liters of water.

Spray bath:

Prepare a mixture with 1.5 mL of Ticoff with 1 liter of water and sprinkle with fine drop to the whole animal.

Facilities:

Apply 3 mL of Ticoff per liter of water.

RETIREMENT PERIOD

Immersion baths:

Meat animals: 3 days

Milk: 6 hours.

PRECAUTIONS

Moderately toxic product. Do not transport or store with food, clothing or fodder. Keep out of the reach of children and pets. Do not store at home room. This product should not be exposed or handled by pregnant women, lactating women and children under 18 years of age. Do not reuse the container, break up and shut up.

CONSULT THE VETERINARY PHYSICIAN