

TRAK

Insecticide and / or acaricide for external use suspension for bath by immersion and aspersion.

REG. RSCO-PEC-INAC-0179-315-008-013

FORMULA

Composition Percentage% IN WEIGHT Amitraz: N¹- (2-4 Dimethyl phenyl) - N - `[[[(2,4-dimethyl phenyl) imino] methyl] -N-methyl-methanimidamide 12.5% (Equivalent to 125 g / L)

Inert ingredients 87.5%

Total 100%



PRESENTATIONS

100 mL, 1 and 8L.

INDICATIONS

TRAK is effective to control the following ectoparasites:

- *Boophilus annulatus*
- *Boophilus microplus*
- *Amblyomma habraeum*
- *Amblyomma americanum*
- *Amblyomma cajennense*
- *Amblyomma maculatum*
- *Damalinia bovis*
- *Haematopinus, spp.*
- *Solenoptes capilatus*
- *Linognathus vituli* *Trichodectes canis*
- *Linognathus setosus*
- *Demodex canis*
- *Otodectes cynotis*
- *Dermacentor albipictus*
- *Rhipicephalus appendiculatus*
- *Otobius megnini*
- *Sarcoptes, spp.*
- *Psoroptes communis*
- *Chorioptes bovis*

INSTRUCTIONS FOR USE

ALWAYS CALIBER YOUR APPLICATION TEAM

SPRAYING AGAINST MITTENS IN CATTLE

TRAK should be diluted in a proportion of 20 mL. For every 10 lts. of water. Use at least 4 liters of the prepared solution for a bovine of approximately 300 kg.

SPRAYING AGAINST SARNA AND LICE IN PIGS

TRAK should be diluted in a proportion of 40 mL. For every 10 lts. of water. Use at least 2 liters of the solution prepared for a pig of approximately 100 kg. Do two treatments with intervals of 7 - 10 days between the first and the second

FLOWING IN FACILITIES

TRAK should be used in a proportion of 80 ml. For every 10 lts. of water

IMMERSION AGAINST MUSHROOMS IN CATTLE

Load: Add 1.6 lts. of TRAK and 6 kg. Of stabilizer of bath simultaneously for each 1000 liters of water.

Recharge: To readjust the level of the bath, 3.2 liters of TRAK and 12 kg should be added. of bath stabilizer simultaneously for every 1000 liters of water.

NOTE: When the bathroom loaded with TRAK is not being used, add 6 kg. of stabilizer per 1000 liters of water every two months. It is recommended to pre-premix in a TRAK pre-mixer basin and water basin as well as the preservative before adding the product to the bath. How to avoid the formation of lumps of the conservator in the bathroom. The pH of the bath should be determined after adding the preservative. The pH should be 12 - 15. If not, you should add more conservative.

For the control of the tick of one or more hosts, the strategic treatment must be carried out depending on the degree of infestation, tick species and geographic location of the animals.

TRAK can be used in cattle, sheep, goats, pigs and dogs, except miniature breeds and puppies under three months of age. TRAK should not be used on horses.

The meat of animals treated with TRAK can be consumed 24 hrs after starting the treatment.

Milk obtained from animals treated with TRAK can be consumed the same day of treatment.

WARNINGS

Its use in equines is not recommended

Do not store next to the food.

Keep out of reach of children

Avoid discarding containers in tanks, waterways and ditches, as it is harmful to aquatic organisms.

TRAK can be used in cattle, sheep, goats, pigs and dogs, except miniature breeds and puppies under three months of age

CONSULT THE VETERINARY PHYSICIAN