

PROVIDEAN VIRATEC 10 CV 4L

Vaccine against Distemper, Parvovirus, Hepatitis, Respiratory Disease (by Adenovirus type II), Parainfluenza, Coronavirus and Leptospirosis of the Canines

REG. SAGARPA B-2083-029

FORMULA

Each dose of 1 mL reconstituted contains attenuated viruses (50% DICC / dose):

Canine parvovirus ($\geq 10^{6.5}$), Distemper ($\geq 10^3$), Canine Adenovirus Type II ($\geq 10^{2.5}$), Canine Parainfluenza ($\geq 10^5$). Inactivated canine coronavirus virus ($\geq 10^5$ TCID₅₀ / dose)

Bacteria of *Leptospira interrogans* serovar *icterohaemorrhagiae* ($\geq 10^7$ Bacteria / dose), *Leptospira interrogans* serovar *canicola* ($\geq 10^7$ Bacteria / dose), *Leptospira interrogans* serovar *Pomona* ($\geq 10^7$ Bacteria / dose), and *Leptospira interrogans* serovar *grippotyphosa* ($\geq 10^7$ Bacteria / dose).

PRESENTATIONS

10 bottles 1 freeze-dried dose each
10 bottles with diluent

INDICATIONS

Vaccination of dogs in good general health is recommended, with proper nutritional development, deworming, without febrile status and out of stress due to adverse environmental conditions. Primary vaccination: In puppies from 6 weeks of age apply a three-dose vaccination program, maintaining an interval of 3 weeks in each of them. It is advisable to have finished the vaccination program before 12 weeks of age to induce the window period of exposure of the puppies during their socialization phase. Revaccination: Apply a booster dose every year.

DOSAGE AND ADMINISTRATION

Canines: 1 mL

Using a needle and sterile syringe, aseptically extract the diluent fraction and reconstitute the lyophilized fraction. Stir until a homogeneous solution is obtained. Administer 1 mL intramuscularly or subcutaneously.

WARNINGS

Keep out of the reach of children. Keep the product in its original container between 2° and 8° C. Do not freeze. Use all the content Use sterile disposable syringes for their application. Avoid vaccination of pregnant females. All vaccines can cause anaphylactic reactions, in which case apply corticosteroid.

VETERINARY USE

CONSULT THE VETERINARY DOCTOR

Made in Argentina by:

Tecnovax S.A, Luis Viale 2835 (C1416DNU) Ciudad Autónoma de Bs. As, Argentina.

SENASA: Certificado N° 06-147. Est. Elaborador N° 8520.

Tel./Fax: (+54) (11) 4585-9695 /4899-0568.

www.tecnovax.com

D.T (Director Científico): Dra. Marcela Iglesias M.P 7140.

