PROVIDEAN VIRATEC 6CV®

Vaccine against Distemper, Parvovirus, Hepatitis, Respiratory Disease (by Adenovirus type II), Parainfluenza and Coronavirus of the Canines

REG. SAGARPA B-2083-026

FORMULA

Each dose of 1 mL reconstituted contains attenuated virus (TCID 50% / dose): Canine Parvovirus (≥106.5), Distemper (≥103), Canine Adenovirus Type II (≥102.5), Canine Parainfluenza (≥105). Coronavirus inactivated virus. Sterile diluent c.s.p 1 mL

PRESENTATIONS

10 bottles 1 freeze-dried dose each 10 bottles with diluent

INDICATIONS

Vaccination of dogs in good general health is recommended, with proper nutritional development, deworming, without febrile status and out of stress due to adverse environmental conditions.

Primary vaccination: In puppies from 6 weeks of age apply a three-dose vaccination program, maintaining an interval of 3 weeks in each of them. It is advisable to have finished the vaccination program before 12 weeks of age to induce the window period of exposure of the puppies during their socialization phase.

Revaccination: Apply a booster dose every year.

DOSAGE AND ADMINISTRATION

Canines: 1 mL

Using a needle and sterile syringe, aseptically extract the diluent fraction and reconstitute the lyophilized fraction. Stir until a homogeneous solution is obtained. Administer 1 ml intramuscularly or subcutaneously.

WARNINGS

Keep out of the reach of children. Keep the product in its original container between 2° and 8° C. Do not freeze. Use all the contents immediately after rehydrating. Use sterile disposable syringes for their application. Avoid vaccination of pregnant females. All vaccines can cause anaphylactic reactions, in which case apply corticosteroid.

VETERINARY USE CONSULT THE VETERINARY DOCTOR

Made in Argentina by: Tecnovax S.A, Luis Viale 2835 (C1416DNU) Ciudad Autónoma de Bs. As, Argentina. SENASA: Certificado N° 06-147. Est. Elaborador N° 8520. Tel./Fax: (+54) (11) 4585-9695 /4899-0568. www.tecnovax.com D.T (Director Científico): Dra. Marcela Iglesias M.P 7140.

