

ZIMACAL®

Reg. SAGARPA Q-2083-164

FORMULA

Each 100 mL contains:

Magnesium 987 mg

Calcium 795 mg

Zinc 670 mg

Vehicle, c.b.p. 100 mL

PRESENTATIONS

Bottles with 500 mL



INDICATIONS

Coadjuvant for the regeneration of the hoof in cases of gabarro, laminitis, abscesses, ulcers and plantar fissures.

Coadjuvant for the treatment of hypocalcemia, slow uterine involution and weak postpartum animals. For the prevention and treatment of tetany of grasses, in paddocks with green shoots after rains with low concentrations of Magnesium and Zinc.

Treatment of metabolic imbalances due to nutritional deficiencies of Magnesium and Zinc. Coadjuvant to reduce the period of adaptation in animals entering the pen.

USE IN

Bovine, ovine and porcine.

DOSE

Bovine, ovine and swine:

Therapeutic: 7 mL per 100 Kg of weight. Preventive: 5 mL per 100 Kg of weight. Frequency of application:

Therapeutic: every 10 days until remission of the signs and no more than 3 repetitions, following the preventive dose.

Preventive: Every 30 days.

ROUTE OF ADMINISTRATION

Only subcutaneous

WARNINGS

In advance consult with the Nutritionist the total contribution of Zinc, Magnesium and Calcium in the diet to determine the dose and time to be supplemented in a parenteral way.

Withdrawal time: 0 days.

Shake it before using.

Keep out of the reach of children. Store in a cool, dry place, protected from light.

CONSULT THE VETERINARY PHYSICIAN VETERINARY USE