

"Agricultural use"



Thiabendazole

Agricultural Fungicide / Wettable Powder

**PERCENTAGE COMPOSITION**

**Active ingredient:**

Thiabendazole: 2- (thiazol-4-yl) benzimidazole  
(Equivalent to 600 g ai / kg at 20 °C)

**Inert ingredients:**

Wetting agent, Dispersants, Thinner

**Total**

Reg: RSCO-FUNG-0399-0254-002-60

**% in weigh**

**60.0**

**40.0**

**100.0**



"Harmful if inhaled"

Registered, developed and distributed by:

**LAPISA, SA DE CV**

Carr. La Piedad-Guadalajara Km. 5.5  
Camelinas Col., CP 59375, La Piedad,  
Michoacan, Mexico. Tel. : 01 (352) 526 13 00.

[www.lapisa.com](http://www.lapisa.com)

MADE IN CHINA AND / OR IN MEXICO

"This material is not a label, the function of this document is illustrative."

**CAUTION**

## "STOP: READ THE LABEL BEFORE USING THE PRODUCT"

### "PRECAUTIONS AND WARNINGS FOR USE"

Use the appropriate and complete protective equipment necessary to safely perform handling operations for the preparation, the loading of the application equipment and its application: cotton cap and overalls, dust mask, goggles or chemical type lenses, gloves and neoprene boots. Keep stored in a safe place.

### "PRECAUTIONS DURING PRODUCT HANDLING"

"DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT".

"DO NOT TRANSPORT OR STORE ALONG WITH FOOD PRODUCTS, CLOTHES OR FORAGE."

"KEEP OUT OF REACH OF CHILDREN, PREGNANT, LACTATION AND ANIMAL WOMEN DOMESTICOS ". "WASH YOUR HANDS AFTER USING THE PRODUCT AND BEFORE CONSUMING FOOD".

"DO NOT STORE IN HOUSES ROOM". "DO NOT USE THE PRODUCT WITHOUT THE PROPER PROTECTION EQUIPMENT".

"PERSONS UNDER 18 YEARS OLD SHOULD NOT HANDLE THIS PRODUCT".

"AT THE END OF THE WORK DAY BATH AND PUT CLEAN CLOTHES".

"DO NOT INGIRE, INHALE AND AVOID CONTACT WITH EYES AND SKIN".

"DO NOT MIX OR PREPARE THE PRODUCT WITH YOUR HANDS, USE GLOVES, USE A PIECE OF WOOD OR OTHER APPROPRIATE MATERIAL ".

"DO NOT UNLOCK THE NOZZLE WITH THE MOUTH".

### "IN CASE OF INTOXICATION, TAKE THE PATIENT WITH THE DOCTOR AND SHOW THIS LABEL"

#### "FIRST AID"

In case of contact with skin or mouth, wash immediately.

Remove the subject from the contaminated area. Remove contaminated clothing and wash exposed parts with plenty of water. If vomiting occurs, flip the subject over the side and ensure that you breathe without difficulty.

### "RECOMMENDATIONS TO THE DOCTOR"

**Chemical Group:** Benzimidazole.

**Antidote:** In case of ingestion of the product, a gastric lavage should be performed. Wash your mouth with water and administer activated carbon mixed with water (1 g per kg of weight). Do not administer milk or anything that contains grease.

**Signs and symptoms of poisoning:** In case of ingestion or inhalation may have side effects Gastrointestinal, transient effects of nausea, vomiting, weakness, drowsiness, dizziness and headache.

**Treatment:** Use symptomatic treatment.

### "MEASURES FOR PROTECTION TO THE ENVIRONMENT"

"DURING THE HANDLING OF THE PRODUCT, AVOID THE CONTAMINATION OF SOILS, RIVERS, LAGOONS, ARROYOS, DAMS, CHANNELS OR WATER DEPOSITS, NOT WASHING OR WASTEING IN THE PESTICIDE RESIDUES OR EMPTY CONTAINERS".

"MANAGE THE EMPTY CONTAINER AND THE WASTE OF THE PRODUCT AS PROVIDED IN GENERAL LAW FOR THE PREVENTION AND INTEGRAL MANAGEMENT OF WASTE, ITS REGULATIONS OR THE MANAGEMENT PLAN FOR EMPTY CONTAINERS OF PESTICIDES, REGISTERED BEFORE THE SEMARNAT ".

"IMPROPER USE OF THIS PRODUCT MAY CONTAMINATE UNDERGROUND WATER. AVOID HANDLING NEAR WATER WELLS AND DO NOT APPLY WHERE THE LEVEL OF THE AQUIFER MANTES IS LITTLE DEEP (75 CM DEPTH) AND THE SOILS ARE VERY PERMEABLE (SANDY) ".

"IN THE EVENT OF ACCIDENTAL FALL OF THE PRODUCT, PERSONAL PROTECTIVE EQUIPMENT SHOULD BE USED, RECOVER THE PRODUCT THAT IS USEFUL UNTIL WHERE POSSIBLE, AND OTHERWISE COLLECT IT IN A HERMETIC CONTAINER AND TAKE IT TO THE AUTHORIZED HAZARDOUS WASTE DISPOSAL CENTER MORE NEAR".



**"Use EXCLUSIVELY IN CROP AND PEST HERE RECOMMENDED"**

**"INSTRUCTIONS FOR USE"**

**"Always calibrate application equipment"**

CULTURE	PLAGUE	DOSAGE / ha
dad (0)	the black scab dad <i>Rhizoctonia solani</i>	2-3kg Seed application the groove bottom, and two applications "in drench" at 60 and 90 days after emergence.
Lemon Lime Tangerine Grapefruit Orange Tangerine Chayote	anthracnose ( <i>Colletotrichum gloeosporioides</i> )	0.5- 0.7 kg / ha Foliar applications perform two with an interval of 14 days, with a volume of 620-720 L of water per ha.
Squash Squash Pepino Melon Watermelon Strawberry Zarzamora	root rot ( <i>Pythium sp.</i> , <i>Sclerotinia sp.</i> , <i>Rhizoctonia sp.</i> , <i>Fusarium sp.</i> And / or <i>Verticillium sp.</i> )	2.0- 3.0 kg / ha Perform one drench application of 10 to 20 days after plant emergence. Application volume of 800 L water / ha, 15 mL of the solution per plant.
	Anthracnose ( <i>Colletotrichum acutatum</i> )	1.5- 2.0 kg / Ha Make two applications foliage 7-day interval with a volume of water of 260-360 l / ha.
Papaya	Anthracnose ( <i>Colletotrichum gloeosporioides</i> )	1.0- 2.0 g / L water Make two applications to the foliage and / or the bank of the fruit with interval of 7 days, with a volume of water of 530-630 l of water / ha.
Avocado (fifteen)	Anthracnose ( <i>Colletotrichum gloeosporioides</i> )	0.75- 1.5 kg Perform two foliar applications directed to the foliage and fruit with interval of 7 days between each.

( ) Guard interval in days between the last application and harvest.

TIME ZONES reentry TREATED:

DO NOT ENTER THE AREA TREATED BEFORE 12 HRS.

METHOD TO PREPARE AND IMPLEMENT THE PRODUCT

Carefully open the bag using the mask.

Calibrate the application equipment, in a separate container place a small amount of water, add the product at the recommended dose on the label and stirring, then empty this premix tank and adding the entire water calculated calibration. It is recommended to maintain stirring.

CONTRAINDICATIONS

Do not apply with strong winds or temperatures above 35 ° C.

PHITOTOXICITY

This product is not phytotoxic to crops here indicated if applied at the recommended dose on the label.

INCOMPATIBILITY

This product must not be mixed with other products.

"RESISTANCE MANAGEMENT"

"TO PREVENT THE DEVELOPMENT POPULATIONS RESISTANT, ALWAYS OBSERVE THE DOSE AND FREQUENCY OF APPLICATION; AVOID REPEATED USE OF THIS PRODUCT, alternating with other groups CHEMICAL different modes of action and different mechanisms of detoxification and by supporting other control methods "

WARRANTY: LAPISA SA DE CV, ensures product composition as specified in this label.

However, under no circumstances liable for misuse of the product or damage that could be caused by it.

